

**Eldorado Fire at Walker Ranch  
Vegetation Reestablishment Monitoring  
2002/2007  
Boulder, County  
Colorado**

Prepared for:  
Boulder County Parks and Open Space  
Small Grants Program 2001  
P.O. Box 471  
Boulder, CO 80503

December 10, 2007

Prepared by:



**Patrick Murphy  
1554 North Street  
Boulder, Colorado 80304  
303-444-4358**

## TABLE OF CONTENTS

<b>Eldorado Fire at Walker Ranch,</b> .....	<b>1</b>
<b>Vegetation Reestablishment Monitoring</b> .....	<b>1</b>
Boulder, County Colorado .....	1
<b>Abstract</b> .....	<b>1</b>
<b>Summary</b> .....	<b>2</b>
<b>Introduction</b> .....	<b>4</b>
Seeding .....	5
Mulching .....	6
Contour Straw Wattles (a.k.a. straw logs).....	6
Contour Log Felling .....	7
<b>Methods</b> .....	<b>10</b>
Sample Site Selection and Documentation .....	10
Data collection.....	10
Vegetation Cover Sampling .....	10
Climate Data.....	13
Classification.....	13
Cover classes and pseudospecies .....	13
Ordination.....	14
<b>Results &amp; Discussion:</b> .....	<b>17</b>
Classification.....	17
Ordination.....	19
Group Descriptions – Synthesis of 2002 Classification and Ordination with vegetation comparison to 2007 .....	29
Climatic Factors .....	54
<b>References</b> .....	<b>58</b>
<b>Appendices</b> .....	<b>59</b>
Appendix 1. Cover Data Tables .....	59
Appendix 2. Species Importance in 2002 and 2007 based on 18 samples.....	72
Appendix 3. Photographs .....	76
Appendix 4. Eldorado Fire, Walker Ranch TWINSPAN 2-Way Table with Sample and Species Classification for 2002 Data.....	95
Appendix 5. 2002 Species & Site Attribute Data.....	96
Appendix 6. Computer files and GPS Sample Coordinates.....	121

## LIST OF FIGURES

Figure 1. Eldorado Fire at Walker Ranch aerial photography with; burn perimeter, burn severity mapping, treatment areas, and sample locations. Coordinate grid is UTM NAD 27 meter. ....	8
Figure 2. Eldorado Fire at Walker Ranch USGS topographic map with; burn perimeter, burn severity mapping, treatment areas, and sample locations. Coordinate grid is UTM NAD 27 meter. ....	9
Figure 3. Sample location markers .....	11
Figure 4. Point-intercept sampling device and the point-intercept transect layout.....	12
Figure 5. Frequency distribution of cover values. ....	14
Figure 6. TWINSpan classification of samples based on 2002 vegetation cover data.....	21
Figure 7. Axes 1-2 – Environmental/treatment/growth form vectors with sample sites (2002).....	23
Figure 8. Axes 1-2 Sites, classification groups and vectors (2002).....	24
Figure 9. Axes 1-2 Selected Species (2002). ....	25
Figure 10. Axes 1-2 Species, groups and environmental vectors (2002). ....	26
Figure 11. Axes 1-2-3, Sites, and Treatments (2002). ....	27
Figure 12. Axes 1-2-3, Sites, and Environmental Factors (2002). ....	28
Figure 13a. Ground cover of the TWINSpan Classification Groups with Growth Form Composition (2002).....	29
Figure 13b. Ground cover of the TWINSpan Classification Groups with Growth Form Composition (2007).....	30
Figure 14. Boulder Annual Precipitation 1893-October 2007.....	54
Figure 15. Boulder Climate Diagrams for 2000 – October 2007. (continued next page).....	56
Figure 15. Boulder Climate Diagrams for 2000 – October 2007. (concluded) .....	57

## LIST OF TABLES

Table 1. Summary of sample locations, treatments, burn severity, and topographic characteristics. ..	10
Table 2. Summary of Site Factors used in Ordination.....	16
Table 3. The 31 most important 2002 species based on cover and frequency in all 18 samples.....	18
Table 4. CANOCO Correlation Matrix for 2002 Data. ....	22
Table 5. Precipitation deviation from the mean 2000 to 2006.....	55

# **Eldorado Fire at Walker Ranch, Vegetation Reestablishment Monitoring**

Boulder, County Colorado

## **Abstract**

This study quantitatively monitored vegetation recovery after fire at 18 locations that were recorded with GPS, marked with survey caps and photographed to allow long-term analysis. The purpose of this study was to provide data that would quantitatively describe post-fire and post-treatment conditions, and monitor change over time. No attempt has been made to compare post-fire vegetation with pre-fire vegetation since no quantitative data were collected prior to the fire. Evaluation of reclamation treatment effectiveness was also beyond the scope of this study since statistically adequate sampling would require a much larger sample size that would include the presence of controls. Controls would require that areas in needed of treatment, not be treated. A complete evaluation of the effectiveness of reclamation would also require a more exact assessment of the vegetation response to burn intensity as well as pre-fire vegetation conditions.

Sites were subjectively selected in 2002 based primarily on post-burn treatments in addition to variation in topography and burn intensity. The vegetation cover data in 2002 and the current 2007 data were collected with a point-intercept scope that allows the integration of forest canopy cover with ground cover, by allowing a single sample point to be projected both upward and downward. This methodology allowed the incorporation of the cover provided by standing dead trees as well as the incorporation of future changes in forest canopy as live tree species become reestablished. The data were also recorded in a manner that allows vegetation and ground cover under the tree canopy to be distinguished from the same values in open areas between trees.

Species composition and species dominance were measured using a combination of a 100 square meter plot for species density, and point-intercept sampling (100 sample points) on a 50-meter transect to quantify species cover dominance. This same methodology has been used by City of Boulder Open Space to monitor tall grass prairie and prairie dog habitat.

The recovery of these burned areas needed to be monitored in order to answer the following questions:

1. How well did the erosion control and seeding work?
2. Which reclamation species did best?
3. Which reclamation species were not successful?
4. How well did the natives recover on their own?
5. Are noxious weeds becoming a problem, and if so which species?
6. What is the current risk of erosion due to bare soil?

## Summary

### 1. How well did the erosion control and seeding work?

The purpose of erosion control is to minimize rill and gully formation. This can best be accomplished by minimizing the amount of bare ground exposed to raindrop impacts and overland sheet flow.

The average bare soil in the 16 burned samples decreased about 44% between 2002 and 2007.

Bare Soil	Average	St.Dev.	Low/High	Sample 17 unburned	Sample 18 unburned
2002	36%	9%	16% - 54%	33%	1%
2007	16%	8.3%	3% - 31%	26%	1%

The average vegetation cover in the 16 burned samples increased about 48% from 2002 to 2007.

Vegetation Cover	Average	St.Dev.	Low/High	Sample 17 unburned	Sample 18 unburned
2002	30%	6%	21% - 39%	26%	63%
2007	44.4%	11.3%	28% - 62%	42%	58%

The Revised Universal Soil Loss Equation RUSLE indicates that when all other factors are held constant (e.g., slope, soil texture), a vegetation cover of about 30% results in the greatest proportional reduction in erosion.

The classification subgroup that most consistently received seeding (Group B n=6) had an average vegetation cover of 27% in 2002 and 32.2% in 2007. About 13% vegetation cover was provided by the reclamation species in 2002 and about 2% was still present in 2007. The 2002 report made the following comment, "... the question should be asked whether or not it would be more reasonable to simply increase ground cover by 13% using persistent mulch (e.g. wood fiber/chips) and allow natives to recover." This may be true, but there was also a reduced amount of cheatgrass observed in the seeded samples in 2007. This is an interesting correlation, but why this might be happening is unknown.

### 2. Which reclamation species did best?

Within the classification subgroup that most consistently received seeding (Group B n=6), mountain brome (*Ceratochloa carinata*) provide an average of 5.4% (range 0% – 10%) cover in 2002 and 0.0% in 2007, slender wheatgrass (*Elymus trachycaulus*) provided about 5.4% (range 0% – 11%) in 2002 and about 1.7% (range 0% - 7%) in 2007, regreen (*Triticum aestivum x Elytrigia elongata*) provided about 1% (range 0% - 3%) in 2002 and 0.0% in 2007, and blue grama (*Chondrosium gracile*) provided about 0.8% (range 0% - 3%) in 2002 and 0.33% (range 0% - 2%) in 2007.

### 3. Which reclamation species were not successful?

Blue grama and regreen did not seem to provide significant cover over the two-year period following the fire (2002) and all of the species have decreased since 2002. However, there may be a relationship between areas that were seeded and a reduced amount of cheatgrass in 2007.

### 4. How well did the natives recover on their own?

Within the classification subgroup that was severely burned but not reseeded, and received only contour log installation (Group A n=3), the average vegetation cover was 37% (s.d. = 2%, range 35% - 39%) in 2002 and 53.3% (s.d. = 4.9%, range 50% - 59%) in 2007. Although this seems much more successful than the seeded Group B (see question 1), Group A (cover increased from 27% to 32%) and B (cover increased from 37% to 53.3%) were on different types of sites. Group A was on steeper NE facing slopes with more trees, and Group B was on flatter NW facing slopes (see Figure 8). This along with other undefined factors may be responsible for the difference.

5. Are noxious weeds becoming a problem, and if so which species?

The predominant weedy species in 2002 were mullein (*Verbascum thapsus*) with 1.28% cover and 78% frequency, Jim hill mustard (*Sisymbrium altissimum*) with 1.5% cover and 61% frequency, cheatgrass (*Anisantha tectorum*) with 1% cover and 72% frequency, Canada thistle (*Breea arvensis*) with 0.56% cover and 39% frequency, and alyssum (*Alyssum minus*) with 0.33% cover and 28% frequency. The predominant weedy species in 2007 were cheatgrass with a 10 fold increase in cover to about 10% with 78% frequency. The cheatgrass problem was greatest in Groups C (17.4%) and D (22%), and least in Groups A (4.7%), B (2%), and E (0%). Group B was the group that most consistently received the seeding treatment. All of the other weedy species have reduced cover with the exception of tumble knapweed (*Acosta diffusa*) which has increased from 0% cover and 11% frequency to 0.5 % cover with 11% frequency. In both years the knapweed was found in samples 9 and 11.

The total cover of introduced species has increased somewhat over the last 5 years. In 2002 the total cover of introduced species was 8% (using the cover value that ignores overstory) and has increased to 12% based primarily on the increase in cheatgrass since most of the other weed species have decreased in cover. When these cover values are compared to the total vegetation cover at the sample locations, the relative cover of the introduced species averaged 25% in 2002 and 28% in 2007. The interesting weed, tobacco weed (*Nicotiana attenuata*) that was observed in 2002 was not found in 2007.

6. What is the current risk of erosion due to bare soil?

Based on a subjective assessment that includes familiarity with the RUSLE annual erosion prediction model, the risk is low. Vegetation cover in the burn areas (Samples 1-16) averaged 30% (range 21% - 39%) in 2002 and 44% (range 28% - 62%) in 2007. The overall ground cover (includes vegetation, litter, rock and standing dead vegetation) averaged 58% (range 43% - 82%) in 2002 and 83.6% (range 68% - 97%) in 2007.

## Introduction

The Eldorado fire at Walker Ranch produced areas with variable impacts related to vegetation and soil exposure. Intensive reclamation efforts of selected areas have helped to reduce the risk of severe erosion and the introduction of non-native noxious weedy species. At the same time there is a risk that disturbance activities related to reclamation, including the use of introduced species in the reclamation mix, may have negative effects on long-term native vegetation recovery. This study compares current (2007) conditions to the baseline (2002) conditions and is an essential component of “adaptive management”. The actual success or failure of these reclamation methods related to vegetation establishment and reduction of erodible bare soil was not previously monitored.

Adaptive management necessitates a critical review of management actions in order to refine and adjust management practices based on actual results. The purpose of this study was to establish vegetation monitoring sites that will provide long-term quantitative data on vegetation establishment and relate the results to environmental conditions such as slope, aspect, climate, severity of burn, and post-burn treatment. Results include sample specific summary of vegetation conditions, classification of all samples based on the year 2002 vegetation characteristics, and ordination of all samples in the environmental/treatment gradient to allow some assessment of results based on site conditions and treatment.

This study was not designed to provide statistical assessment of success or failure of the reclamation methods. The hypothesis testing that statistical assessment requires, mandates an intensive sampling methodology that must include sample adequacy determination, and untreated controls. The current study utilizes numerical analysis techniques to determine trends that point the way for future hypothesis testing if that is ultimately desired. As stated in the original proposal the following questions were addressed.

The recovery of these burned areas needs to be monitored to answer such questions as:

1. How well did the erosion control and seeding work?
2. Which species did best?
3. Which were not successful?
4. How well did the natives recover on their own?
5. Are weeds becoming a problem, and if so which species?
6. What is the current risk of erosion due to bare soil?

Figures 1(aerial photography) & 2(USGS topography) provide maps of the burn area and the approximate treatment zones with eighteen sample locations. The perimeter of the burn area is identified with a red line. The burn severity areas are identified with black perimeter lines around a red hatch pattern for severely burned areas and a blue hatch pattern for moderately burned areas. The areas within the red perimeter line but outside of the hatch areas were typically unburned but were sometimes lightly burned or had received small spot burns. The treatment areas are identified by magenta perimeter lines and the treatment associated with each area is described in the Methods section of this report. The actual reclamation treatments are

described below. Note that climate stations shown in Figures 1 and 2 were not reestablished in 2007.

The following details regarding the burn were provided by Boulder Area Sustainability Information Network (BASIN) and can be found at the following web site:

<http://bcn.boulder.co.us/basin/news/Eldorado.html>

The Walker Ranch Fire, also referred to as the Eldorado Fire, began about 2 p.m. Friday, Sept. 15, 2000 and consumed some 1100 acres before it was completely contained on Wednesday, September 20th. There was no loss of life and no structures were burned. The Boulder Daily Camera reports that more than 500 people, 74 fire-fighting engines, 273,000 gallons of water and 133,000 gallons of fire retardant were used to bring the fire under control.

The burn occurred along and near South Boulder Creek west and north of Eldorado Springs State Park, primarily on Boulder County Open Space land, but in after the fire there were potentially impacts to the drinking water supplies of the cities of Denver, Louisville and Lafayette which all draw source water for their treatment facilities from South Boulder Creek.

The Colorado State Forest Service has provided a more detailed graphic presentation of the burn chronology at <http://lamar.colostate.edu/~csfsbo/fire.htm#>

Boulder County provided the following details on the burn impacts and reclamation efforts at the following web site. [http://www.co.boulder.co.us/openspace/resources/ecology/walker\\_fire.htm](http://www.co.boulder.co.us/openspace/resources/ecology/walker_fire.htm)  
Of the 1,062 acres that were burned in the Eldorado Fire, 450 acres were moderately or severely burned.

275 acres were severely burned.

- 50% to 100% of the canopy was burned
- Needles are gone
- Ground cover was partially consumed
- Weak areas of hydrophobic or water resistant soils may be present

175 acres were moderately burned.

- Up to 50% of the canopy was burned
- Needles are gone from many trees, but not all
- Ground cover was partially consumed
- There may be weak areas of hydrophobic soils may be present.

## Seeding

Minimal seeding will be used since this fire was relatively small and there are ample native seed sources surrounding the burned areas. Seed will only be hand broadcast onto areas that are at high risk for severe erosion or noxious weed invasion. The following seed mix is being used:

- 28% of the mix will be Blue grama (*Bouteloua gracilis*), a dominant native grass, (Authors Note: the actual value included 30% and 32% in site specific mixes Claire DeLeo – Eldorado Area Rehabilitation Plan Revisions and Summary, undated BCPOS).
- 25% will be Mountain brome (*Bromus marginatus*), a pioneer native grass,

(Authors Note: the actual value included 27% in site specific mixes Claire DeLeo – Eldorado Area Rehabilitation Plan Revisions and Summary, undated BCPOS).

- 32% will be Slender wheatgrass (*Elymus trachycaulus*), a short-lived native perennial, and,

(Authors Note: the actual value included 35% and 37% in site specific mixes Claire DeLeo – Eldorado Area Rehabilitation Plan Revisions and Summary, undated BCPOS).

- 15% will be "Regreen", a sterile hybrid of Cereal wheat (*Triticum aestivum*) and Tall wheatgrass (*Elytrigia elongata*), short-lived perennials.

(Authors Note: the actual value included 4% and 10% in site specific mixes Claire DeLeo – Eldorado Area Rehabilitation Plan Revisions and Summary, undated BCPOS).

(Current Author's Comment: Although blue grama grass is a local native, it is not typically the dominant in the upper foothill areas of this burn. Mountain brome (a.k.a. *Ceratochloa carinata*) is an introduced species in Colorado although native in some states of the U.S. This species did not naturally occur in the burn areas prior to planting. Slender wheatgrass is also a local native species, but was probably not abundant in the burn area prior to seeding.)

The reasons why introduced or non-local species are used are based on the facts that native species, especially local native species, are either unavailable or expensive. Not all areas were seeded. The areas that were seeded were selected because they were the greatest concern with regard to erosion or noxious weed invasion. The non-native species were chosen based on their ability to produce quick ground cover and yet be short-lived and allow the reestablishment of native species.

(Current Author's Comments: This study allows some evaluation of whether these species were successful at providing quick cover, and permit evaluation of whether or not they are short-lived.)

### **Mulching**

Mulching reduces the erosive action of raindrops hitting bare soil and overland sheet flow. Certified weed-free winter wheat straw is applied at 1 ton (about 50 bales) per acre. Seeds remaining in the straw will germinate and provide a temporary ground cover until native plants can reestablish.

Mulching is also used in conjunction with seeding to provide a protective cover for seeds by reducing soil moisture evaporation.

### **Contour Straw Wattles (a.k.a. straw logs)**

Straw wattles are used on severe to moderately burned slopes with less than 30 percent of the original ground cover remaining. They increase infiltration, add roughness, reduce erosion, and help retain eroded soil on slopes. They are also used to supplement erosion control in areas that do not have enough large trees for contour log felling and in rocky areas where contour log felling is difficult to implement.

Straw wattles are cylinders of compressed weed-free straw. They are made of either wheat or rice straw, and are 8 to 12 inches in diameter and 20 to 25 feet long. They are encased in jute, nylon, or other bio/photo-degradable materials. When installed on the contour of a slope they form a continuous barrier that intercepts water and sediment running down the slope. Straw wattles are effective for about 3 years.


### **Contour Log Felling**

When the original ground cover is lost during a fire, the soil is at risk for erosion. Drainage ways may flood more frequently from increased runoff on the burned slopes. Contour log felling can reduce erosion from rainwater that runs down a slope by cutting dead trees so they fall perpendicular to the main direction of the slope. This technique is used on burned slopes where about 50% or more of the tree canopy is destroyed.

Sawyers cut trees, dropping the trunks along the contour of the slope leaving stumps about 12 inches high to brace the tree from sliding downhill. Tree limbs are removed so that the log lies flat on the ground. Soil is then packed under the log to slow the flow of water and facilitate the deposition of sediment on the upslope side of the log.

A discussion of potential water quality impact was also provided by the following BASIN web site: <http://bcn.boulder.co.us/basin/forum/walkerWQ.html>

By: Donna Scott, City of Boulder, Water Quality and Environmental Services Potential water quality impacts involve a major tributary to Boulder Creek and a drinking water supply reservoir. It is estimated that 500,000 people receive their drinking water from water resources affected by this fire. These include South Boulder Creek, which is a drinking water source for the cities of Louisville, Lafayette and the town of Superior and Gross Reservoir, a water supply for Denver Water as well as serving Arvada. In addition, several small drainage ways cross the area. Tom Davis Gulch is an intermittent stream which runs west to east right through the most heavily burned areas and is a tributary to South Boulder Creek, just upstream of the city of Lafayette's and the town of Louisville's diversion structures. Gross Reservoir's northern corner is within a few hundred feet of the fire area, and South Boulder Creek forms most of the eastern boundary of the fire.



**Figure 1. Eldorado Fire at Walker Ranch aerial photography with; burn perimeter, burn severity mapping, treatment areas, and sample locations. Coordinate grid is UTM NAD 27 meter.**




Figure 2 provides a map of the burn area and the approximate treatment zones with eighteen sample locations. The perimeter of the burn area is identified with a red line. The burn severity areas are identified with black perimeter lines with a red hatch pattern for severely burned areas and a blue hatch pattern for moderately burned areas. The areas within the red perimeter line but outside of the hatch areas were typically unburned but were sometimes lightly burned or had received small spot burns. The treatments areas are identified by magenta perimeter lines and the treatment associated with each area is described in the Methods section of this report.

**Figure 2. Eldorado Fire at Walker Ranch USGS topographic map with; burn perimeter, burn severity mapping, treatment areas, and sample locations. Coordinate grid is UTM NAD 27 meter.**

## Methods

### Sample Site Selection and Documentation

Eighteen samples were subjectively selected from the study area (Figures 1 & 2) and stratified as much as possible to include the range of burn severity, reclamation treatments and topographic position. Sample 18 was selected from a site that was outside of the burn perimeter in a densely forested site. The mapping of the treatment areas was not precise and required on-site adjustment of plot location and orientation to best fulfill the targeted combination of treatments that each sample was intended to represent. Table 1 summarizes the results of sample selection. The sample transects were marked by a large survey cap and brown carsonite post at the start point, and a white fiberglass pole and small aluminum caps at the end point (Figure 3). The start and end points were recorded with a Trimble Geoplotter 3 GPS unit that was accurate to approximately +/- 1.5 meters.

**Table 1. Summary of sample locations, treatments, burn severity, and topographic characteristics.**

Sample ID	Treatment Unit	Contour	Straw logs (acres)	Seed (acres)	Mulch (acres)	Treatment Summary	Burn Class	Slope-Aspect	
	A1	X	3.1	6	6	Everything	Severe	M - NE	M = moderate
1	A2	X	2	3	5	Everything	Severe	M - NE	
2	F1	X	10.1		2.5	C& S-Logs & Mulch	Severe	M - SE	
3	F2	X	10.1		4.5	C& S-Logs & Mulch	Severe	S - SE	S = Steep
4	I	X	5.8		4	C& S-Logs & Mulch	Severe	M - SE	
5	L	X	5.7		8	C& S-Logs & Mulch	Severe	M - NE	
6	D2	X		5	4	C& Seed & Mulch	Severe	S - W	
7	E	X	2.9			C& Straw Logs	Moderate	Ridge - SE	
	J	X	1.9			C& Straw Logs	Severe		
8 & 9	G1	X		8		C& Seed only	Severe	S - NW	Top & bottom of slope
10	C2	X		some		C& some seed	Severe	Ridge - ENE	Seeded top
11	SBCTrail			1.5 some		Seed only	Moderate	S - SW	
12 & 13	B	X				Contour only	Severe	S - ENE	Top & bottom of slope
	B2	X				Contour only	Moderate		
14	C	X				Contour only	Severe	Drain - ENE	
	D1	X				Contour only	Severe		
	G2	X				Contour only	Severe+Mod		
	H	X				Contour only	Light+Severe		
	K	X				Contour only	Moderate + Sev.		
15	Other Burn					None	Severe	S - NE	
16	Other Burn					None	Severe	S - NW	
17	No burn interior					None	None	M - W	
18	No burn dense exterior					None	None	S - NNE	
<b>18</b>	<b>TOTAL Samples</b>								

### Data collection

#### **Vegetation Cover Sampling**

Vegetation cover was originally sampled at the 18 transects (Figures 1 & 2) on July 16-19, 2002. The 2007 sampling occurred July 21-24. Each 50 meter transect was sampled with 100 points using a point-intercept optical device (Figure 4). Two points were sampled at each meter, one on either side of the transect at 0.5 meter from the transect centerline (Figure 4). The point-intercept optical device uses high quality optics and cross-hairs to project a point in an upward as well as downward direction, allowing the canopy of living or dead trees to be recorded. Whenever an upward point recorded a hit on tree canopy



Sample Transect Start Point



Sample Transect End Point




Aluminum Cap with Transect Number

### Figure 3. Sample location markers

(either alive or dead), the additional downward hits were recorded separately to allow the discrimination of points that occurred under a tree canopy.

Each sample point recorded first-hit (top canopy) and additional hits for vegetation by species, as well as litter, bare soil, rock, and standing-dead vegetation. Litter was defined as dead organic matter in contact with the soil or within 1 cm. of the soil surface. “Standing-dead” was defined as organic matter that was over one year old (i.e., not this year’s vegetation growth) that was not yet in contact with soil and was below about 1.5 meter above the ground. In this study, all dead standing trees were recorded by species and kept distinct from “standing dead” organic matter.



**Figure 4. Point-intercept sampling device and the point-intercept transect layout.**

Species within one meter (3.28 ft.) of the transect centerline were also recorded as "present". This allowed species with low cover to be represented in the data and provided a species density per 100 square meters (i.e., 50 meters long by 2 meters wide plot). The transect was subjectively oriented to best represent the community/treatment target. This cover sampling methodology is identical to the vegetation monitoring used by Boulder City Open Space and Mountain Parks in their prairie dog studies, and the Boulder County study at the Doniphan Property prairie dog revegetation site.

In 2002 photographs were only taken at the start points of the transects. A horizontally oriented and a vertically photograph were taken at each sample. In 2007 horizontally oriented photographs were taken at both ends of the transects, and an occasional vertical photograph was also taken.

## Climate Data

The climate data were not collected in 2007. A climate diagram was prepared that presents the monthly temperature, precipitation, and potential evapotranspiration for the last 5 years and compares it to the last 100 years of record.

## Classification

The vegetation cover data were classified in 2002 using the TWINSpan (Hill, 1979) divisive classification program with the strict convergence criteria of Oksanen & Minchin (1997). This program defines groups of samples based on the similarity of their species composition, and simultaneously groups the species that tend to occur together within each classification group. Certain settings must be entered for TWINSpan to perform the classification. These settings are described in the following section.

## Cover classes and pseudospecies

Because TWINSpan was originally designed to be used for Presence-Absence (i.e. frequency) data rather than quantitative (i.e. abundance) data, the use of "Pseudospecies" was incorporated into the computer program. The concept of pseudospecies allows greater weight to be given to higher quantitative values. The first setting of TWINSpan for pseudospecies requires that the cover classes (i.e. cut levels) be defined. TWINSpan allows a maximum of nine cut levels.

In this study the cut levels were defined as 0.0, 0.2, 1.1, 3.1, 5.1, 7.1, 9.1, 11.1, >11.1. The first cut level included all species cover values that were greater than zero, cut level two included all values that were greater than or equal to 0.2%, cut level three included all values greater than or equal to 1.1%, etc. The 0.2 cut level was used to distinguish those species that were present, but were not tallied as a hit by the cover sample. All of these "present" species were given a value of 0.1. Almost all species had cover values less than 19%. Figure 5 shows the frequency distribution of cover values at the targeted cut levels. A species that was present (SPEX) in the plot but not "hit" was assigned a value of 0.1 and was assigned one pseudospecies (i.e. SPEX1). A species with a cover value of 10% would be assigned eight pseudospecies (i.e. SPEX1, SPEX2, SPEX3, SPEX4, SPEX5, SPE6, SPE7, SPE8) because it could be found to occur within eight of the nine possible cover classes. If the cover classes are not equal in size to each other, there is a *de facto* weighting of the data. For example the species with 10% cover would occur in eight out of nine possible classes, so its weighted cover would be 89% (i.e. 8/9). The net effect is to positively weight the lower cover values and negatively weight the higher cover values. This can be compensated by the weighting option of the TWINSpan program. For this study the weights given to each class were respectively; 9,000, 45,000, 60,000, 67,500, 90,000, 105,000, 115,714, 123,750, 190,000. For example, the species with 10% cover would be given a weight of 123,750 and would be multiplied by 89 (i.e., the  $8/9 = 89\%$  value) = 11,013,750. The maximum weighted value for a species with 100% cover would be  $100 \times 1,000,000 = 100,000,000$ . The net weighting for the 10% species is  $11,013,750/100,000,000 = 11\%$ . The final result is that the cover classes are "unweighted", and more representative of the true cover values.

The TWINSpan results were used to define the sample and species associations that have resulted from the fire disturbance and reclamation efforts. This is presented in a dendrogram that includes those species, growth forms (i.e., introduced annual forbs, native perennial grasses, etc.), and selected environmental factors that were most closely associated with each division. The determination of these associated growth forms and environmental factors was accomplished using DISCRIM variant of the TWINSpan program (ter Braak 1982).

### Ordination

The samples were ordinated using CANOCO Version 4.0 (ter Braak 1999) which incorporates the strict convergence criteria of Oksanen & Minchin (1997). The ordinations of samples, species, and environmental vectors were produced separately for different combinations of two axes, as well as a simulated three-dimensional scatter plot. The “environmental factors” included reclamation treatments, independent site factors, vegetation related site factors, and growth forms. Table 2 summarizes the environmental factors used in the ordination. Slope was measured in percent, and aspect was composed of the combination of “easterliness” (sine of aspect) and “northerliness” (cosine of aspect). Aspect was recorded with respect to true north. Abbreviations used throughout this report are as follows:

#### Treatments


- Contour = Contour log felling
- Strwlogs = Straw logs (wattles)
- Seed = Application of seed
- Mulch = Application of Mulch
- Burn Intensity
- Burn = Burn Intensity

#### Independent Site Factors

- Sloppcnt = Slope in percent
- AspETN = Easterly component of aspect relative to true north
- AspNTN = Northerly component of aspect relative to true north

#### Vegetation Related Site Factors

- StndDead = Standing Dead vegetation that has been dead for at least one full year, and that is not in contact with mineral soil, litter, or the ground surface.
- Litter = Non-living organic matter, that is in contact with the mineral soil, other litter or the ground surface.



**Figure 5. Frequency distribution of cover values.**

Baresoil = Soil with no vegetation or litter cover.

Rock = Rock that is greater than 1cm in size in at least one dimension.

TotVeg = Total vegetation cover

SpeDen = Species density (i.e., number of species found within the 100sq.m. [2m x 50m] transect)

Grndcov = Ground cover that will help reduce erosion. Includes total vegetation cover, rock, litter, and standing dead.

#### Growth Forms

IAF = Introduced Annual & Biennial Forbs

IAG = Introduced Annual Grasses

IPF = Introduced Perennial Forbs

IPGC = Introduced Perennial Grasses (cool season)

NAF = Native Annual & Biennial Forbs

NPF = Native Perennial Forbs

NPGC = Native Perennial Grasses (cool season)

NPGW = Native Perennial Grasses (warm season)

S = Native Shrubs

T = Native Trees

F = Native Ferns

M = Moss

The associations as defined by the TWINSPAN classification were also represented within the ordination using connecting lines or distinct symbols to show the distribution of the classification groups within the ordination. The species that were found to distinguish the classification groups were also presented in separate attribute plots. The attribute plots show the actual cover values of the selected species for each sample. The cover values are represented by circles with a diameter that is proportional to the cover value, and the circles are centered on the location of the sample in the ordination. It is then possible to get a realistic idea of the distribution of cover values for each species with respect to the plant associations and the environmental factors.

**Table 2. Summary of Site Factors used in Ordination.**

Ordination Site Factors	Samples																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
<b>Treatments</b>																		
Contour	1	1	1	1	1	1	1	1	1	1		1	1	1				
Strwlogs	1	1	1	1	1		1											
Seed	1					1		1	1	1	1							
Mulch	1	1	1	1	1	1												
<b>Burn Intensity</b>																		
Burn	4	4	4	4	4	4	3	4	4	4	3	4	4	4	4	4	0	0
<b>Independent Site Factors</b>																		
Sloppcnt	18%	29%	29%	27%	29%	40%	18%	32%	32%	11%	32%	36%	45%	25%	40%	25%	36%	32%
AspETN	0.2924	0.2250	0.3907	0.7986	0.8746	-0.9563	-0.1908	-0.2924	-0.4540	0.9986	0.6820	0.9962	0.9063	0.9925	0.9205	-0.7547	-0.9563	0.6820
AspNTN	0.9563	-0.9744	-0.9205	-0.6018	0.4848	0.2924	-0.9816	0.9563	0.8910	-0.0523	-0.7314	0.0872	0.4226	0.1219	0.3907	0.6561	0.2924	0.7314
<b>Vegetation Related Site Factors</b>																		
StndDead	2	1	4	2	3	0	0	0	1	3	0	1	0	0	3	0	2	0
Litter	43	19	30	26	30	19	20	16	17	15	19	13	9	15	16	20	29	79
Baresoil	18	49	37	29	31	49	43	44	51	57	55	37	42	41	31	54	33	2
Rock	0	1	1	5	1	7	3	9	6	0	0	10	7	1	8	1	10	0
TotVeg	41	31	32	40	37	24	36	31	26	26	31	51	46	53	45	30	28	73
SpeDen	49	43	47	46	36	36	39	46	39	35	44	32	31	48	57	40	39	25
Grndcov	86	52	67	73	71	50	59	56	50	44	50	75	62	69	72	51	69	152
<b>Growth Forms</b>																		
NAF	0.1	2.0	1.0	2.0	0.1	1.0	3.0	1.0	1.0	3.0	5.0	11.0	4.0	18.0	1.1	6.0	2.0	0.1
IAF	4.0	1.0	10.0	10.0	20.0	1.0	0.1	0.1	0.1	0.1	5.0	6.0	0.1	4.0	11.0	1.0	1.0	0.0
IAG	1.0	1.0	0.1	7.0	7.0	1.1	0.1	0.1	0.0	0.0	0.1	0.0	0.0	6.0	0.1	0.1	0.1	0.0
NPF	1.0	20.0	3.0	9.0	7.0	10.0	18.0	5.0	13.0	5.0	3.0	14.0	21.0	14.0	13.0	23.0	7.0	0.1
IPF	1.1	0.1	6.0	1.0	0.1	1.0	1.0	0.1	0.0	0.1	1.0	4.0	0.1	0.1	0.1	0.1	0.1	0.1
NPGC	18	6	7	10	2	7	11	10	3	5	11	1	4	2	15	0.1	6	12
IPGC	12	0.1	0.1	0.1	1	3	1	11	7	10	2	0	0	0	0	0	0	0
NPGW	3.0	0.0	0.0	0.1	0.1	0.1	0.0	3.0	0.1	1.0	1.0	0.0	0.0	0.0	1.0	0.0	3.0	0.0
Shrub	1.0	1.0	5.0	1.0	0.1	0.1	2.0	1.0	1.0	2.0	3.0	12.0	13.0	2.0	4.0	0.1	9.0	0.1
Tree	9.0	12.0	5.0	1.0	7.1	11.0	6.0	10.0	11.0	4.0	29.0	13.0	13.0	18.0	16.0	9.0	1.0	64.0
Fern	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Moss	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0

## Results & Discussion:

### Data Tables

The vegetation cover data for 2002 and 2007 are presented in Appendix 1 as summary tables that include the original cover data, average and relative cover for each species and growth form, total cover for each sample, and the number of species that occurred within the 100 sq.m. plot for each sample.

This results and discussion section will repeat the findings from 2002 that describe the “baseline” conditions and how those were used to initially classify the samples into groups. The new information will simply compare the overall, group, and individual sample changes over time.

Table 3 presents the 31 most important species observed in 2002, based on a combination of constancy (a term that is equivalent to frequency when comparing samples) and total cover. This list includes those species that were non-preferentials as well as some species that TWINSPAN determined were indicator species for some of the groups. The complete list of the 2002 species used for this analysis sorted by relative importance is presented in Appendix 2 along with the 2007 list of most important species. Also in Appendix 2 is a list of the species with the greatest increase or decrease in importance since the 2002 sampling. A listing of those species that were newly observed as well as those that were observed in 2002 but not in 2007 is included in Appendix 2. The photographs of samples are presented in Appendix 3.

### Classification

The TWINSPAN 2-way classification table is presented in Appendix 4. This table presents both the sample and species divisions produced by the TWINSPAN program. The TWINSPAN classification summary is presented in Figure 6 and defined five groups (A-E) based on species composition. The species listed at the top of Figure 6 were common in all groups, and were non-discriminatory. Because samples were selected primarily from fire disturbance sites, no attempt has been made to classify the communities based on typical undisturbed plant community structure.

This classification integrates the results of burn intensity, site conditions, reclamation treatment, and natural revegetation. Figure 6 demonstrates that much of the grouping based on species composition correlates well with reclamation treatments. NO simplistic or definitive conclusion can be made, however, regarding the cause and effect of the treatments, since the treatments were applied subjectively in areas of greatest need. For example, Group A (Samples 12, 13, 14) was severely burned with no treatment and was found to be typified by the presence of quaking aspen (*Populus tremuloides*). The relative absence of aspen in all of the other “treatment” groups does not mean that “treatment” excluded regeneration of aspen. It is more likely that aspen occurred in areas that were more likely to regenerate on their own, or were less of an erosion risk, and were therefore not selected to receive reclamation treatment. On the other hand, when the treatment included seeding, especially with non-native species, the change in community composition due to these species can be assumed to be caused by reclamation efforts.

Although five groups were defined by the classification, only four occur within the burn area. A description of the groups follows the next section, which describes the Ordination results.

**Table 3. The 31 most important 2002 species based on cover and frequency in all 18 samples.**

Scientific Name	Relative Importance (%)	Constancy (%)	Average Cover All-Hits (%)
Pinus ponderosa ssp. scopulorum (dead)	100.00	66.67	5.61
*Carex pensylvanica ssp. heliophila	65.92	77.78	3.17
⊙Ceanothus fendleri	53.28	94.44	2.11
⊙Phacelia heterophylla	53.24	77.78	2.56
Pseudotsuga menziesii (dead)	42.93	55.56	2.89
⊙Verbascum thapsus	36.36	94.44	1.44
Sisymbrium altissimum	28.10	61.11	1.72
⊙Chenopodium simplex	27.66	77.78	1.33
*Anisantha tectorum	24.72	72.22	1.28
Elymus trachycaulus	19.34	33.33	2.17
*Ceratochloa carinata	18.36	33.33	2.06
⊙Penstemon virens	15.92	88.89	0.67
Artemisia ludoviciana	12.94	72.22	0.67
Geranium caespitosum ssp. caespitosum	12.69	77.78	0.61
Carex spp.	11.10	50.00	0.83
Corydalis aurea	9.95	55.56	0.67
Artemisia frigida	8.96	50.00	0.67
⊙Campanula rotundifolia	8.11	77.78	0.39
⊙Chenopodium leptophyllum	8.11	77.78	0.39
Grindelia squarrosa	7.25	44.44	0.61
Triticum aestivum x Elytrigia elongata	7.19	61.11	0.44
Helianthus pumilus	6.95	66.67	0.39
Breaa arvensis	6.34	38.89	0.61
Apocynum androsaemifolium	6.30	22.22	1.06
Chondrosium gracile	5.94	44.44	0.50
Epilobium brachycarpum	5.88	66.67	0.33
Leucopoa kingii	5.88	66.67	0.33
⊙Astragalus miser var. oblongifolius	5.41	72.22	0.28
*Populus tremuloides	4.93	22.22	0.83
Pseudotsuga menziesii	4.88	5.56	3.28
Physocarpus monogynus	4.63	44.44	0.39

⊙ = Nonpreferential species. This symbol marks those species that are evenly distributed among all of the classification groups. The other species showed some degree of preference for specific groups.

\* = Indicator species determined by TWINSpan. These species were closely associated (high fidelity) with specific groups and little or no association with other groups. Although some indicator species were also relatively dominant in the groups, an indicator species may also be present with low cover values. Since the indicator typically occurs only in a specific group and not in the others it is a good indicator for group affiliation when it is observed in a sample.

## Ordination

The purpose of the ordination graphic is to reduce an extremely complex system to a fewer number of manageable factors. No presumption is made that all of the most important factors are represented. This is a first approximation that should be refined in an iterative process. The cross-tabulated correlation matrix of the ordination axes and environmental factors is presented in Table 4. Significant correlations are highlighted in blue ( $P < .05$ ) or red ( $P < .01$ ).

CANOCO provides ordinations with four axes that are typically represented two axes at a time. The fourth axis was found to be well represented by the other three axes and is not presented here. These axes do not represent a specific environmental/treatment factor, but rather a two step process that maximizes the dispersion of the samples on each axis. The first step of the ordination “arranges” the samples based on vegetation similarities. The second step then further refines the “arrangement” by creating axes that are linear combinations of all of the environmental/treatment factors. The vectors associated with the environmental/treatment factors are “best fit” axes for these specific factors. The direction of the vector from the origin indicates the direction of increasing values for the factor, and the length of the vector indicates the relative importance of the factor (when compared to the other measured factors) for explaining the variability found in the data.

The ordination graphics are presented in four two-dimensional figures using axes 1 & 2 (i.e., the x- and y-axes respectively) from the CANOCO results, and two simulated 3-dimensional figures. Most of the variability in the data were represented by the first two axes but Samples 17 and 18, the unburned samples, were found to be separated from the cluster of other samples along the 3<sup>rd</sup> axis. Sample 18, the dense forest sample that was outside of the burn area, was excluded from the ordination because it was so distinctive that it forced all of the other samples into a tight cluster. Sample 17 although distinctive, had similar species composition and allowed a reasonable ordination result.

Figure 7 presents the sample sites with respect to the “environmental” vectors. The treatment vectors are in red, the environmental site vectors are in black, and the growth form vectors are in green. The direction of the vectors indicates the direction of increasing value for that factor. The length of the vector indicates the relative importance of that vector in the ordination. The position and magnitude of the environmental vector is determined by its ability to maximize its contribution to an explanation of the species and site ordinations. The position of a sample site in the ordination is determined both by its species composition, and its environmental site factors.

Figure 8 presents the samples with respect to the environmental vectors and the zones that are occupied by the groups defined by the TWINSPLAN classification results.

Figure 9 presents those species that had the greatest effect on the ordination results. Those species with red font were indicators of the classification groups. The position of a species in the ordination represents its center of distribution and is determined by its cover distribution among the sample sites.

Figure 10 combines the three previous figures and allows simultaneous comparison of samples, species, and environmental factors.

Figure 11 is the 3-dimensional representation of Figure 7 with the samples and treatment vectors projected to permit some visualization of different paired combinations of the three axes. The floor of the simulated cube is a representation of axes 1 & 2, the right wall represents axes 2 & 3, and the left back wall represents axes 1 & 3. The most significant item to notice is that although sample 17 appears to be

close to the other samples when only axes 1 & 2 are used, it can be seen to be separated from the other samples along the 3<sup>rd</sup> axis.

Figure 12 is similar to Figure 11 but includes the environmental factors.

The graphics in Appendix 5 plot the actual cover values for species at the sample locations in order to allow a more accurate representation of the distribution of species within this same ordination.


## Eldorado Fire, Walker Ranch TWINSPAN Results

### 18 samples

Species found in all Groups (non-preferentials) and in >70% of the samples.

- Ceanothus fendleri (native shrub NS)  
 Penstemon virens (native perennial forb NPF)  
 Phacelia heterophylla (NPF)  
 Chenopodium leptophyllum (native annual forb NAF)  
 Chenopodium simplex (NAF)  
 Campanula rotundifolia (NPF)  
 Astragalus miser (NPF)

**Verbascum thapsus (Introduced Biennial Forb)**



Text in red indicates non-native species.

• This symbol in magenta is used to mark the indicator species identified by TWINSPAN

• This symbol in blue is used to mark the indicator environmental/treatment/growth form factors that are associated with the classification levels. The program DISCRIM was used to identify these indicators. Additional factors that were “preferentials” are sometimes listed below the indicators.

The number of samples in each division is at the top of each column.

Single parenthesis = (xxxxx) = center in a moderately wide distribution

Double parenthesis = ((xxxxx)) = center in a broad distribution

Bracketed parenthesis = [(xxxxx)] = secondary center in moderate distribution


**This classification dendrogram presents the species, samples and environmental/treatment/growth form factors that are associated with the groups defined by the TWINSPAN program based on the vegetation composition of the samples.**

**The program DESCRIM was used to determine the environmental/treatment/growth form factors that are associated with the classification groups that were originally determined by TWINSPAN based on vegetation composition.**

TWINSPAN Classification results using total hit (i.e., first and additional hits) data.

**Figure 6. TWINSPAN classification of samples based on 2002 vegetation cover data**





**Figure 7. Axes 1-2 – Environmental/treatment/growth form vectors with sample sites (2002).**

The treatment vectors are in red, the “environmental” site vectors are in black, and the growth form vectors are in green. The direction of the vectors indicates the direction of increasing value for that factor. The length of the vector indicates the relative importance of that vector in the ordination.




Figure 8. Axes 1-2 Sites, classification groups and vectors (2002).




Figure 9. Axes 1-2 Selected Species (2002).




Figure 10. Axes 1-2 Species, groups and environmental vectors (2002).




Figure 11. Axes 1-2-3, Sites, and Treatments (2002).





Figure 12. Axes 1-2-3, Sites, and Environmental Factors (2002).

Group Descriptions – Synthesis of 2002 Classification and Ordination with vegetation comparison to 2007

The Groups were originally defined in 2002. Most of the following information for 2002 is schematically summarized in Figure 6 (TWINSPAN classification dendrogram) and Figure 48a (Ground Cover of the TWINSPAN classification groups). The 2007 Ground Cover for the Groups is presented in Figure 48b.

Although five groups were defined by the classification, only four occur within the burn area. Group E was a single sample in an unburned forest. A side-by-side description of the groups follows the 2002 and 2007 figures, and a side-by-side comparison of samples follows the group comparisons.




IAF = Introduced Annual & Biennial Forbs  
 IAG = Introduced Annual Grasses  
 IPF = Introduced Perennial Forbs  
 IPGC = Introduced Perennial Grasses (cool season)

NAF = Native Annual & Biennial Forbs  
 NPF = Native Perennial Forbs  
 NPGC = Native Perennial Grasses (cool season)  
 NPGW = Native Perennial Grasses (warm season)

S = Native Shrubs  
 T = Native Trees  
 F = Native Ferns  
 M = Moss

**Figure 13a. Ground cover of the TWINSPAN Classification Groups with Growth Form Composition (2002).**




IAF = Introduced Annual & Biennial Forbs  
 IAG = Introduced Annual Grasses  
 IPF = Introduced Perennial Forbs  
 IPGC = Introduced Perennial Grasses (cool season)

NAF = Native Annual & Biennial Forbs  
 NPGC = Native Perennial Grasses (cool season)  
 NPGW = Native Perennial Grasses (warm season)  
 NPF = Native Perennial Forbs

S = Native Shrubs  
 T = Native Trees  
 F = Native Ferns  
 M = Moss



**Figure 13b. Ground cover of the TWINSpan Classification Groups with Growth Form Composition (2007).**

The following are side by side comparisons of the ground cover and growth forms for 2002 and 2007.





Group **A** in 2002 was composed of 3 samples (12, 13, 14) that were defined in 2002 by the presence of the indicator species quaking aspen. All three samples received the same severe burn classification and received only contour log installation. Group A was distinguished by higher total vegetation cover (36.5%) that was composed of only a small percentage of introduced species (4.3% see Figure 48). It also tended to have higher cover values of native annual forbs and native shrubs. Species density averaged 37.0 species/100 sq.m. in 2002 and 38 species/100 sq.m. in 2007.

The percentage of bare ground was reduced from 36.5% to 8.0% over the five years and the vegetation cover increased from 37% to 53.3%. The native annual forbs decreased from 7.7% to 1.3%, but the introduced annual grass cheatgrass increased from 1% to 4.7%. A large increase in shrub cover from 7% to 18.7% is noteworthy and was primarily buckbrush (*Ceanothus fendleri*) with 14.3% cover. The dead standing trees provided 10% cover in 2002 but reduced to 2.3% in 2007. This phenomenon was repeated in most groups as the dead standing trees fell over and were then evaluated as “standing dead” cover. Note that the standing dead cover increased from 0% to 6% over the 5 years. “Standing dead” is defined as organic matter that is over one year old (i.e., not this year’s vegetation growth) that is not yet in contact with soil and is below about 1.5 meter above the ground. In this study, all dead standing trees were recorded by species and kept distinct from “standing dead” organic matter. This group and Group B (which was seeded) had the least problem with cheatgrass.





Group **B** in 2002 was composed of 6 samples (1, 6, 8, 9, 10, 16) that were defined primarily by having been severely burned and seeded. Mountain brome and slender wheatgrass were the predominantly distinctive species for this group with the exception of Sample 16. Sample 16 was not seeded but was included in this group primarily due to the standing dead ponderosa and Douglas fir. Sample 16 could be considered transitional to Group A. This group also typically had a more northerly aspect, and more bare soil. These stands may have been denser stands with a denser layer of duff and fewer understory species prior to the fire. The fire may have burned hotter here, and the combination of these factors leads to the decision to apply seed along with some other combination of treatments. The sites may have appeared to be more sterile resulting in a post-fire management decision to apply seed. Total vegetation cover was 26.5% with 7% composed of introduced species and an additional 5.3% provided by slender wheatgrass (one of the reclamation species that is a non-local native). Species density averaged 40.8 species/100 sq.m. in 2002 and 38.2 species/100 sq.m. in 2007.

Bare soil decreased from 40.7% in 2002 to 20.0% in 2007. Vegetation cover increased somewhat from 26.5% to 32.2%, but the greatest increase in ground cover was litter which increased from 20.3% to 35.5%. The dead standing trees have converted to “standing dead” cover as described for Group A. A large decrease in the introduced perennial grasses was primarily a decrease in the reclamation species. The native perennial reclamation grass slender wheatgrass (which isn't a local native species) also decreased, but was compensated by increases in the native grasses such as thickspike wheatgrass (*Elymus lanceolatus*) and sun sedge (*Carex pensylvanica* ssp. *heliophila*). This was more than compensated by the increase in native perennial forbs and shrubs (primarily buckbrush). This group which was seeded and Group A had the least problem with cheatgrass.



Group **C** in 2002 was composed of 5 samples (2, 4, 5, 7, 11) that were defined primarily by having been severely burned, with contour log felling, straw logs, and mulch treatment with no seeding. Samples 7 and 11 were rated as moderately burned and sample 7 received no mulch or seed, and site 11 received only seeding. A suite of species also typified this group, such as Jim Hill mustard (*Sisymbrium altissimum*), hairy golden aster (*Heterotheca villosa*), and wild buckwheat (*Eriogonum umbellatum* var. *umbellatum*). The abundance of standing dead ponderosa pine combined with sedge (*Carex pensylvanica* ssp. *heliophila*) also typified this group. This group had the highest percentage cover of introduced species with about 9.6% cover, but most of this cover was provided by annual introduced species (2.8% from cheatgrass, 5% from Jim Hill mustard, 1.4% from alyssum). Although most of these sites received no seeding, there was still a trace of the introduced reclamation grass species in these areas. This may have been due to migration of seed from the seeded areas due to either human or natural causes such as surface water flow mobilization of the seed. Species density averaged 41.6 species/100 sq.m. in 2002 and 29.2 species/100 sq.m. in 2007. This reduction was due primarily to fewer native perennial forb species.

Bare soil decreased from 33.4% in 2002 to 17.8% in 2007. Vegetation cover increased from 30.5% to 47.2%. Litter increased slightly from 22.4% to 27.2%. The dead standing trees have converted to “standing dead” and possibly litter cover as described for Group A. Although native species cover increased from 20.8% to 26%, the introduced weeds increased from 9.6% to 21% primarily due to cheatgrass which had 17.4% cover. The tumble knapweed has also increased from an average cover of <1% to 1.8% but has been restricted to Sample 11 for both 2002 and 2007. The Jim Hill mustard has been reduced from 5% to less than 1%. The reason why cheatgrass is so successful in this group as well as Group D is purely speculation and may simply be coincidence, but I do note that 4 out of 5 samples in Group C received straw logs, and 5 out of 6 of the samples that received straw log treatments have the greatest cheatgrass cover. Also, 4 out of the 6 samples that received the straw log treatment also were mulched. The cheatgrass distribution is probably related to a complex of factors including proximity to human and wildlife trails. The correlation to treatments may simply be due to chance.




Group **D** in 2002 was composed of 3 samples (3, 15, 17) that were relatively intermediate with respect to many site and treatment characteristics, but were defined primarily by the indicator species bladderpod (*Lesquerella montana*), and relatively high values of buckbrush (*Ceanothus fendleri*) similar to Group A. Sample 17 is distinctly separated from the other two samples in the ordination (Figures 11 & 12). Sample 17 may be included in this group due to similar species composition, but may be distinct due to the greater abundance of many of the species because this site was not burned. Sample 17 may be considered a target point on the trajectory of Samples 3 and 15 toward recovery of the more open forested stands in the burn area. Species density averaged 47.7 species/100 sq.m in 2002 and 33 species/100 sq.m. in 2007. This reduction in species density was due primarily to a reduction of native perennial forbs that were present at low levels in 2002.


Bare soil decreased from 31.2% in 2002 to 14.3% in 2007. Vegetation cover increased greatly from 30.5% to 54.7%, but this was due primarily to a large increase in cheatgrass from 0.1% in 2002 to 22% in 2007. Although the native herbaceous cover has decreased slightly from 16.7% to 16.3%, native shrub cover has increased from 4.7% to 14% due primarily to the increase in buckbrush and birchleaf mountain mahogany (*Cercocarpus montanus*). Bladderpod was present in all three samples in 2002 and was an indicator species, but was observed in only one sample in 2007.

This group and Group C had the worst problem with cheatgrass compared to the other groups.

**Sample 18 - 2002**





**Sample 18 - 2007**





Group E was composed of only one off-site sample (Sample 18) in a densely forested north-facing slope. This sample was distinct from all of the other samples because of the dense cover of Douglas fir along with the absence of many of the understory species that were excluded due to the closed canopy. Species density was 25 species/100 sq.m. in 2002 and 38 species in 2007 due primarily to trace amounts of native forbs and grasses.



As might be expected, the overall changes in this undisturbed site were minimal.





Sample 1 - Severe burn, Contour, Straw logs, Seed, Mulch – Group B. Bare ground was greatly reduced and litter greatly increased. Vegetation cover increased only slightly but there was a large reduction in introduced species including the reclamation grasses that was matched by an increase in native forbs and grasses. Cheatgrass here was minimal, as in other seeded samples. Species density has changed from 49 to 45 species/100 sq.m.





Sample 2 - Severe burn, Contour, Straw logs, Mulch – Group C. Bare soil decreased moderately. The dead standing trees reduced greatly and have fallen over providing ground cover as litter. The vegetation cover has increased greatly, but primarily due to cheatgrass. Species density has changed from 43 to 31 species/100 sq.m.





Sample 3 - Severe burn, Contour Straw logs, Mulch – Group D. Bare soil decreased greatly and vegetation cover increased greatly. Although the native species increased moderately, the greatest increase was cheatgrass. Species density has changed from 46 to 28 species/100 sq.m.





Sample 4 - Severe burn, Contour, Straw logs, Mulch – Group C. Bare ground was reduced greatly with an increase in litter. Although native vegetation cover remained about the same, there was a large shift from native grasses to shrubs, primarily buckbrush. The large increase in total vegetation cover was due primarily to cheatgrass. Species density has changed from 46 to 25 species/100 sq.m.





Sample 5 - Severe burn, Contour, Straw logs, Mulch – Group C. Bare soil was greatly reduced, and vegetation cover was increased. Although the site had abundant Jim Hill mustard in 2002, there was none observed in 2007. Cheatgrass and fringed sage (*Artemisia frigid*) are the current site dominants. Species density has changed from 36 to 31 species/100 sq.m.





Sample 6 - Severe burn, Contour, Seed, Mulch – Group B. Bare soil has been reduced, primarily due to a combination of increased vegetation cover and litter. The reclamation grass species appear to have been replaced by native perennial forbs and shrubs. The amount of cheatgrass here was low as in most of the seeded areas. Species density has changed from 35 to 38 species/100 sq.m.





Sample 7 - Moderate burn, Contour, Straw logs – Group C. Most of the dead standing trees have fallen at this site. The bare soil has been reduced and vegetation cover has increased due primarily to the increase in cheatgrass. Golden aster (*Heterotheca foliosa*), fringed sage, and buckbrush are the current dominants. Species density has changed from 38 to 25 species/100 sq.m.





Sample 8 - Severe burn, Contour, Seed – Group B. Bare soil has reduced by about half and litter has increased about the same amount. Some of the dead standing trees have fallen but many are still standing. Reclamation grasses have been replaced by a mix of native perennial forbs and the shrub buckbrush. There was no cheatgrass here as in some of the other seeded areas. Species density has changed from 46 to 32 species/100 sq.m.





Sample 9 - Severe burn, Contour, Seed – Group B. Bare soil has decreased by about half and litter has increased about a third. Some standing dead trees have fallen and accumulated along the transect giving high standing dead cover. Transect was moved due to a trail that can be seen in the 2007 photo. Reclamation grasses have greatly reduced and native perennial forbs dominate. Species density has changed from 39 to 37 species/100 sq.m.





Sample 10 - Severe burn, Contour, Seed – Group B. Bare soil was reduced by more than half and litter increased by more than half. Vegetation cover greatly increased and the reclamation grasses were greatly reduced. The native perennial forbs and shrubs currently dominate. Species density has changed from 35 to 40 species/100 sq.m.





Sample 11 - Moderate burn, Seed – Group C. Bare soil decreased by more than half and the abundant dead standing trees have fallen. Although the vegetation has greatly increased, almost half of the vegetation cover is from weedy species especially cheatgrass and tumble (diffuse) knapweed. Species density has changed from 44 to 34 species/100 sq.m.





Sample 12 - Severe burn, contour only – Group A. Bare soil was greatly reduced and both litter and cover have increased. Some of the dead standing trees have fallen, but many still remain. The greatest increase in vegetation cover was due to the shrub buckbrush. Aspen has actually decreased in cover but still occurs along the transect. Species density has changed from 31 to 34 species/100 sq.m.





Sample 13 - Severe burn, contour only – Group A. Bare soil was greatly reduced and vegetation cover and litter increased. The greatest increase in vegetation cover was due to buckbrush and cheatgrass. Aspen cover has remained stable, but is increasing in stature. Many of the dead standing trees have fallen. Species density has changed from 31 to 39 species/100 sq.m.





Sample 14 - Severe burn, contour only – Group A. Bare ground was reduced to only 3%. Vegetation cover has increased greatly and many of the dead standing trees, or limbs have fallen. *Chenopodium* spp. dominated the cover in 2002, but was not observed in 2007. Aspen is increasing in cover and stature, with an abundance of Agasizz bluegrass (*Poa agasizzensis*). Species density has changed from 48 to 40 species/100 sq.m.





Sample 15 - Severe burn, no treatment – Group D. Start and end points could not be found. New points were installed near the original points using GPS coordinates. Bare soil has decreased greatly and many of the dead standing trees have fallen. Vegetation cover has increased, and buckbrush has increased, but cheatgrass is abundant. Species density has changed from 57 to 32 species/100 sq.m.



Sample 16 - Severe burn, no treatment – Group B. Bare soil has decreased greatly and litter has increased and many of the dead standing trees have fallen. The vegetation cover was dominated by varileaf scorpionweed (*Phacelia heterophila*) in 2002 but is now dominated by fringed sage and cheatgrass. Buckbrush is becoming established. Species density has changed from 40 to 37 species/100 sq.m.



Sample 17 - No burn, no treatment – Group D. This site wasn't burned and is located near a hiking trail. Bare soil and litter have decreased and vegetation cover have increased. Native grasses have reduced but native shrubs have increased. Cheatgrass was present at less than 1% in 2002 but is now present with 10% cover. Species density has changed from 39 to 36 species/100 sq.m.




Sample 18 - No Burn, no treatment – Group E. This unburned site has remained stable. Species density has changed from 24 to 35 species/100 sq.m. The new species were only present in trace amounts.

## Climatic Factors

Although this report does not discuss the historical disturbances or climatic factors that have resulted in the pre-fire plant associations, climatic data for the last 110 years (1894 – 2006) that may be used for that purpose can be found at <http://www.myxyz.org/phmurphy/Download/THRN2006.pdf>. These data for 1894-2007 will be posted in February 2007. The average annual precipitation as well as monthly temperature, precipitation and potential evapotranspiration can provide great insight into the timing and severity of disturbance conditions that existed in the past.

The Figure 14 is the annual precipitation for Boulder from 1894 to October 2007. The mean and +/- one standard deviation is also indicated on the graph as well as polynomial smoothing of the precipitation



**Figure 14. Boulder Annual Precipitation 1893-October 2007.**

It can be seen from Figure 14 that the precipitation was significantly above normal from 1995 to 1999. Table 5 shows that this was followed by below average precipitation in the year of the fire (2000) followed by a year of close to average precipitation in 2001, a very dry year in 2002, followed by above average wet years in 2003 and 2004, with a very wet year in 2004, a below average year in 2005 and a close to average year in 2006. The climate diagram for 2007 found in Figure 15 shows that the January to July period for 2007 was much drier than normal.

**Table 5. Precipitation deviation from the mean 2000 to 2006.**

<b>Year</b>	<b>Precip.</b>	<b>Mean</b>	<b>Deviation</b>
2000	15.96	18.94	(2.98)
2001	18.28	18.94	(0.66)
2002	13.88	18.94	(5.06)
2003	22.02	18.94	3.08
2004	27.17	18.94	8.23
2005	17.52	18.94	(1.42)
2006	19.31	18.94	0.37

The Thornthwaite climate diagrams for the years 2000 to October 2007 are shown in Figure 15. These diagrams show the time and relative intensity of drought periods over time. The periods of potential soil drought occur when potential evapotranspiration (ET) exceeds precipitation. The potential evapotranspiration was calculated according to a modified Thornthwaite formula (Dunne & Leopold 1978) that includes a latitude correction.

When the combined effects of temperature and precipitation for the period of April-May-June are considered, the year 2000 was the 7<sup>th</sup> warmest and driest Spring in the last 110 years, 2001 was the 46<sup>th</sup> (relatively average), and 2002 was the 5<sup>th</sup> warmest and driest Spring. When this same type of comparison is made for the period of April to September; 2000 was the 15<sup>th</sup> warmest and driest Spring/Summer, 2001 was the 36<sup>th</sup>, and 2002 was the 3<sup>rd</sup> warmest and driest in the last 110 years. What this implies is that the year of the reclamation actions (2001) was relatively average, but the second year of growth (2002) was one of the warmest and driest ever. This could have had a significant effect on the subsequent growth responses; however the following two years (2003-2004) were well above average, followed by 2005 that was below average and 2006 that was close to average.




Figure 15. Boulder Climate Diagrams for 2000 – October 2007. (continued next page)





Figure 15. Boulder Climate Diagrams for 2000 – October 2007. (concluded)

## References

- COB, City of Boulder. 1988. Forest Inventory Handbook. City of Boulder Open Space Department and City of Boulder Mountain Parks Division, Parks and Recreation Department.
- Dunne, T.E., Leopold, L.B. 1978. Water in environmental planning. W.H. Freeman and Co., San Francisco. 818 pp.
- Ecotone 2003. Eldorado Fire at Walker Ranch Vegetation Reestablishment Monitoring, Boulder, County Colorado. Report to Boulder County Parks and Open Space Small Grants Program 2001.
- Hill, M.O. 1979. TWINSpan - A FORTRAN program for arranging multivariate data in an ordered two-way table by classification of the individuals and attributes. Microcomputer Power, Ithaca New York, USA.
- Marr, J.W. 1961. Ecosystems of the east slope of the Front Range in Colorado. University of Colorado Studies Series in Biology No. 8. University of Colorado Press.
- Oksanen, J., Minchin, P.R. 1997. Instability of ordination results under changes in input data order: explanations and remedies. *Journal of Vegetation Science* (8) 447-454.
- Shimwell, D.W. 1971. The description and classification of vegetation. University of Washington Press, Seattle. 322 pp.
- ter Braak, C.J.F. 1982. DISCRIM - A modification of TWINSpan (Hill, 1979) to construct simple discriminant functions and to classify attributes, given a hierarchical classification of samples. Institute TNO for Mathematics, Information Processing and Statistics, Wageningen, Netherlands.
- ter Braak, C.J.F. (1987-1997). CANOCO Version 3.15 - A FORTRAN program for canonical community ordination. Microcomputer Power, Ithaca, New York, USA.

# Appendices

## Appendix 1. Cover Data Tables

Combined 2002

Combined 2007

Groups A, B, C, D 2007









## DATA FROM FILE Group A - 2007

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	12D	12DWU	12U	13D	13DWU	13U	14D	14DWU	14U
<b>NATIVE ANNUAL &amp; BIENNIAL FORBS</b>																
<i>Androsace septentrionalis</i>		PYGMYFLOWER ROCKJASMINE	0.00	33.33	0.00	0.00	0.00	P								
<i>Epiobium brachycarpum</i>	EPILOBIUM PANICULATUM	BIGFRUIT WILLOWHERB	0.00	66.67	0.00	0.00	0.00	P			P					
<i>Frasera speciosa</i>		GREEN GENTIAN	0.00	33.33	0.00	0.00	0.00				P					
<i>Grindelia squarrosa</i>		GUMWEED	0.00	100.00	0.00	0.00	0.00	P			P					
<i>Machaeranthera bigelovii</i>		BIGFLOW ASTER	1.33	100.00	2.50	1.33	2.03	4			P			P		
<i>Polygonum douglasii</i>	POLYGONUM SAWATCHENSE,POLYGONUM MONTANUM	DOUGLAS KNOTWEED	0.00	33.33	0.00	0.00	0.00				P					
<b>TOTAL NATIVE ANN. &amp; BIEN. FORBS</b>			1.3	100.0	2.5	1.3	2.1	4	---	---	P	---	---	P	---	---
<b>INTRODUCED ANNUAL &amp; BIENNIAL FORBS</b>																
<i>Alyssum minus</i>		ALYSSUM	0.00	33.33	0.00	0.00	0.00				P					
<i>Carduus nutans ssp. macrolepis</i>		MUSK THISTLE	0.00	33.33	0.00	0.00	0.00							P		
<i>Cynoglossum officinale</i>		HOUND'S TONGUE	0.00	33.33	0.00	0.00	0.00							P		
<i>Lactuca scariola</i>		PRICKLY LETTUCE	0.33	33.33	0.63	0.33	0.51							1		
<i>Sisymbrium altissimum</i>		JIM HILL MUSTARD	0.00	33.33	0.00	0.00	0.00							P		
<i>Tragopogon dubius ssp. major</i>		YELLOW SALSIFY	0.00	66.67	0.00	0.00	0.00	P			P					
<i>Verbascum thapsus</i>		MULLEIN	2.00	100.00	3.75 (3.68)	2.67	4.06	2	(2)		1			3		
<b>TOTAL INTRO. ANN. &amp; BIEN. FORBS</b>			2.3	100.0	4.4	3.0	4.7	2	(2)	---	1	---	---	4	---	---
<b>INTRODUCED ANNUAL GRASSES</b>																
<i>Anisantha tectorum</i>	BROMUS TECTORUM	CHEATGRASS	4.67	66.67	8.75	5.00	7.61				6			8(1)		
<b>TOTAL INTRO. ANN. GRASSES</b>			4.7	66.7	8.8	5.0	7.9	---	---	---	6	---	---	8(1)	---	---
<b>NATIVE PERENNIAL FORBS</b>																
<i>Achillea lanulosa</i>		WESTERN YARROW	3.00	100.00	5.63	3.67	5.58	P			P			9(2)		
<i>Aletes acaulis</i>		STEMLESS INDIAN PARSLEY	0.00	66.67	0.00	0.00	0.00	P			P					
<i>Apocynum androsaemifolium</i>		SPREADING DOGBANE	0.33	66.67	0.63	0.33	0.51	P			1					
<i>Artemisia frigida</i>		FRINGED SAGE	1.33	66.67	2.50	1.33	2.03	2			2					
<i>Artemisia ludoviciana</i>		PASTURE SAGE	0.33	66.67	0.63	0.67	1.02				1(1)			P		
<i>Aster porteri</i>		PORTER'S ASTER	0.33	100.00	0.63	0.33	0.51	1			P			P		
<i>Astragalus laxmannii</i>	ASTRAGALUS ADSURGENS VAR. ROBUSTIER	LAXMANN'S MILKVETCH	0.00	33.33	0.00	0.00	0.00							P		
<i>Astragalus miser var. oblongifolius</i>		WEEDY MILKVETCH	0.00	33.33	0.00	0.00	0.00							P		
<i>Campanula rotundifolia</i>		HAREBELL	0.67	100.00	1.25	0.67	1.02	2			P			P		
<i>Cerastium strictum</i>	CERASTIUM ARVENSE	MOUSE-EAR	0.00	33.33	0.00	0.00	0.00	P								
<i>Drymocallis fissa</i>		POTENTILLA FISSA	0.00	66.67	0.00	0.00	0.00				P			P		
<i>Gaillardia aristata</i>		BLANKETFLOWER	0.00	33.33	0.00	0.00	0.00							P		
<i>Galium septentrionale</i>	GALIUM BOREALE	NORTHERN BEDSTRAW	0.33	66.67	0.63	0.33	0.51				P			1		
<i>Geranium caespitosum ssp. caespitosum</i>		SMALL-LEAF WILD GERANIUM	0.00	66.67	0.00	0.00	0.00	P						P		
<i>Harbouria trachypleura</i>		WHISKBROOM PARSLEY	0.00	66.67	0.00	0.00	0.00	P						P		
<i>Helianthus pumilus</i>		SUNFLOWER	0.33	100.00	0.63	0.33	0.51	P			P			1		
<i>Heliomeris multiflora</i>	GYMNOLOMIA MULTIFLORA	SHOWY GOLDENEYE	0.00	33.33	0.00	0.00	0.00							P		
<i>Heterotheca foliosa</i>		GOLDENASTER	2.33 (2.67)	33.33	4.38 (4.91)	2.67	4.06				7	1		2		
<i>Heterotheca villosa</i>	HETEROOTHECA HORRIDA,CHRYSOPSIS VILLOSA	HAIRY GOLDEN ASTER	3.33 (3.67)	100.00	6.25 (6.75)	3.67	5.58	1			7	1		P		
<i>Lupinus argenteus</i>		SILVER LUPINE	0.00	66.67	0.00	0.00	0.00				P			P		
<i>Monarda fistulosa var. menthifolia</i>		HORSEMINT	0.00	33.33	0.00	0.33	0.51									(1)
<i>Penstemon glaber</i>	PENSTEMON ALPINUS	BEARD TONGUE	0.00	33.33	0.00	0.00	0.00				P					
<i>Penstemon spp.</i>		BEARD-TONGUE	0.00	33.33	0.00	0.00	0.00	P								
<i>Penstemon strictus</i>		ROCKY MOUNTAIN PENSTEMON	0.00	33.33	0.00	0.00	0.00							P		
<i>Penstemon virens</i>		GREEN BEARD-TONGUE	0.00	66.67	0.00	0.00	0.00	P						P		
<i>Phacelia heterophylla</i>		VARIABLE SCORPIONWEED	0.67 (1.00)	66.67	1.25 (1.84)	1.00	1.52	P						2	1	
<i>Scrophularia lancoolata</i>		FIGWORT	0.00	33.33	0.00	0.00	0.00							P		
<i>Solidago missouriensis</i>		MISSOURI GOLDENROD	0.00	33.33	0.00	0.00	0.00							P		
<i>Solidago simplex var. simplex</i>	SOLIDAGO SPATHULATA VAR. NEOMEXICANA	MT. ALBERT GOLDENROD	0.33	100.00	0.63	0.33	0.51	1			P			P		
<b>TOTAL NATIVE PERENNIAL FORBS</b>			13.3 (14.3)	100.0	25.0 (26.4)	15.7	24.7	7	---	---	18(1)	2	---	15(2)	1(1)	---
<b>INTRODUCED PERENNIAL FORBS</b>																
<i>Arabis hirsuta</i>		HAIRY ROCKCRESS	0.00	33.33	0.00	0.00	0.00							P		
<i>Breœa arvensis</i>	CIRSIIUM ARVENSE	CANADA THISTLE	0.67	100.00	1.25	0.67	1.02	1			P			1		
<i>Taraxacum officinale</i>		COMMON DANDELION	0.00	33.33	0.00	0.00	0.00	P								
<b>TOTAL INTRO. PERENNIAL FORBS</b>			0.7	100.0	1.3	0.7	1.1	1	---	---	P	---	---	1	---	---
<b>NATIVE PERENNIAL GRASSES (cool)</b>																
<i>Carex pensylvanica ssp. heliophila</i>	CAREX HELIOPHILA	SUN SEDGE	0.33	33.33	0.63	0.33	0.51							1		
<i>Carex spp.</i>		SEDE	2.33	100.00	4.38	2.33	3.55	4			3			P		
<i>Elymus elymoides</i>	SITANON HYSTRIX	BOTTLEBRUSH SQUIRRELTAIL	0.33	100.00	0.63	0.33	0.51	1			P			P		
<i>Elymus lancoolatus</i>	AGROPYRON LANCEOLATUM,A. DASYSTACHYUM,A. ALBICANS,A. RIP	THICKSPIKE WHEATGRASS	0.67	66.67	1.25	0.67	1.02							2		
<i>Elymus lancoolatus fm. albicans</i>	AGROPYRON DASYSTACHYUM,A. ALBICANS,AGROPYRON RIPARIUM	MONTANA WHEATGRASS	0.00	33.33	0.00	0.00	0.00				P					
<i>Koeleria macrantha</i>	KOELERIA CRISTATA,K. PYRAMIDATA,K. GRAECILIS	JUNE GRASS	0.00	66.67	0.00	0.00	0.00	P			P					
<i>Leucopoa kingii</i>		SPIKE FESCUE	0.33	33.33	0.63	0.33	0.51	1								
<i>Poa agassizensis</i>		AGASSIZ BLUEGRASS	4.00	100.00	7.50 (7.36)	6.33	9.64	2			P			10(5)	(2)	
<b>TOTAL NATIVE PERENNIAL GRASSES (c)</b>			8.0	100.0	15.0	10.3	16.3	8	---	---	3	---	---	13(5)	(2)	---

DATA FROM FILE Group A - 2007

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	12D	12DWU	12U	13D	13DWU	13U	14D	14DWU	14U		
<b>INTRODUCED PERENNIAL GRASSES (cool)</b>																		
<i>Poa compressa</i>		CANADA BLUEGRASS	0.33	66.67	0.63	0.33	0.51	P						1				
<b>TOTAL INTRO. PERENNIAL GRASSES (c)</b>								P	---	---	---	---	---	1	---	---		
<b>NATIVE SHRUBS</b>																		
<i>Ceanothus fendleri</i>		BUCKBRUSH	14.33 (15.00)	100.00	26.88 (27.61)	17.33	26.40	22(2)	2		17(4)			4(1)				
<i>Padus virginiana ssp. melanocarpa</i>	FRUNUS VIRGINIANA SSP. MELANOCARPA	CHOKECHERRY	0.33	66.67	0.63	0.33	0.51	1						P				
<i>Physocarpus monogynus</i>		NINEBARK	1.33	33.33	2.50	1.67	2.54	4(1)										
<i>Ribes cereum</i>		WAX CURRANT	0.33	33.33	0.63	0.33	0.51				1							
<i>Rosa arkansana</i>		ARKANSAS ROSE	0.33	33.33	0.63	0.33	0.51				1							
<i>Rosa woodsii</i>		WOOD'S ROSE	0.33	33.33	0.63	0.33	0.51							1				
<i>Rubus idaeus ssp. melanolasius</i>		AMERICAN RED RASPBERRY	1.00	100.00	1.88	1.33	2.03	1			2(1)			P				
<i>Symphoricarpos rotundifolius</i>	SYMPHORICARPOS OREOPHILUS	MOUNTAIN SNOWBERRY	0.67	33.33	1.25	0.67	1.02							2				
<b>TOTAL NATIVE SHRUBS</b>								28(3)	2	---	21(5)	---	---	7(1)	---	---		
<b>NATIVE TREES</b>																		
<i>Populus tremuloides</i>	POPULUS TREMULA	QUAKING ASPEN	4.00 (3.33)	100.00	7.50 (6.13)	4.67	7.11	P			2			6	2	4		
<i>Pseudotsuga menziesii [dead]</i>		DOUGLAS FIR	2.33 (0.33)	100.00		2.33	3.55			[2]	[1]		[2]			[2]		
<b>TOTAL NATIVE TREES</b>								P	---	---	2	---	---	6	2	4		
Standing dead		STANDING [dead]	6.00 (0.00)	100.00		6.33		[6]			[5]			[7]	[1]			
Litter		LITTER	24.67 (25.33)	100.00		25.33		25			22			27	2			
Bare soil		BARE SOIL	8.00	100.00		8.00		8			13			3				
Rock		ROCK	5.67	100.00		5.67		9			6			2				
<b>TOTALS</b>								92			92			91				
<b>TOTALS (LAYER)</b>								92	2	0	92	2	0	87	5	4		
<b>TOTAL VEGETATION COVER (LAYER)</b>								50(3)	2(2)	0	51(6)	2	0	55(9)	3(3)	4		
<b>TOTAL VEGETATION COVER</b>								53.3 (54.7)			100.0 (100.6)	65.7 (s=0.0)	100.0	50(7)		59(15)		
<b>GROUND COVER (Litter+Rock+Veg+St.Dead)</b>								84(3)	2(2)	0	79(6)	2	0	84(9)	5(3)	4		
<b>SPECIES DENSITY (# of species/100 sq.m.)</b>								34						39		41		
<b>(AVERAGE= 38.0 Std.Dev.= 3.6)</b>																		
<b>SPECIES DENSITY (LAYER)</b>								34	2	0	39	2	0	40	4	1		

(#) = second hit  
 {#} = cover value if tree canopy is excluded  
 [#] = hit on standing dead tree species



DATA FROM FILE Group B

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	01D	06D	06DWU	06U	08D	08DWU	08U	09D	09DWU	09U	10D	16D	16DWU	16U
<b>INTRODUCED PERENNIAL FORBS</b>																					
<i>Brexa arvensis</i>	CIRSIUM ARVENSE	CANADA THISTLE	0.00	66.67	0.00	0.00	0.00	P	P			P			P						
<i>Linaria genistifolia ssp. dalmatica</i>	LINARIA DALMATICA	DALMATIAN TOADFLAX	0.00	16.67	0.00	0.00	0.00												P		
<i>Taraxacum officinale</i>		COMMON DANDELION	0.17	50.00	0.52	0.17	0.41	1				P							P		
<b>TOTAL INTRO. PERENNIAL FORBS</b>			0.2	83.3	0.5	0.2	0.5	1	P	---	---	P	---	---	P	---	---	P	---	---	---
<b>NATIVE PERENNIAL GRASSES (cool)</b>																					
<i>Achnatherum lettermanii</i>	STIPA LETTERMANII	LETTERMAN NEEDLEGRASS	0.00	16.67	0.00	0.00	0.00					P									
<i>Agrostis scabra</i>		TICKLEGRASS	0.00	16.67	0.00	0.00	0.00	P													
<i>Bromopsis lanatipes</i>	BROMUS LANATIPES	WOOLLY BROME	0.17	16.67	0.52	0.17	0.41														1
<i>Carex pensylvanica ssp. heliophila</i>	CAREX HELIOPHILA	SUN SEDGE	1.83	100.00	5.70 (5.56)	2.00	4.98	2	3	(1)		1			3			1	1		
<i>Carex spp.</i>		SEDE	0.33	16.67	1.04	0.33	0.83	2													
<i>Elymus elymoides</i>	SITANION HYSTRIX	BOTTLEBRUSH SQUIRRELTAIL	0.00	33.33	0.00	0.00	0.00	P													P
<i>Elymus lanceolatus</i>	AGROPYRON LANCEOLATUM, A. DASYSTACHYUM, A. ALBICANS, A. RIP	THICKSPIKE WHEATGRASS	3.17	50.00	9.84	3.50	8.71	3(1)	3(1)										8		P
<i>Elymus lanceolatus fm. albicans</i>	AGROPYRON DASYSTACHYUM, A. ALBICANS, AGROPYRON RIPARIUM	MONTANA WHEATGRASS	0.00	33.33	0.00	0.00	0.00														P
<i>Elymus trachycaulus</i>	AGROPYRON TRACHYCAULUM	SLENDER WHEATGRASS	1.67 (1.83)	33.33	5.18 (5.56)	2.17	5.39					7(1)	1		3(1)						
<i>Elymus virginicus</i>		VIRGINIA WILDRYE	0.33	16.67	1.04	0.33	0.83					2									
<i>Festuca arizonica</i>		ARIZONA FESCUE	0.00	16.67	0.00	0.00	0.00														P
<i>Koeleria macrantha</i>	KOELERIA CRISTATA, K. PYRAMIDATA, K. GRACILIS	JUNEGRASS	0.33	83.33	1.04	0.33	0.83	P	P			P			1						1
<i>Leucopoa kingii</i>		SPIKE FESCUE	0.33	66.67	1.04	0.33	0.83	P	P			P									2
<i>Poa agassizensis</i>		AGASSIZ BLUEGRASS	0.00	33.33	0.00	0.17	0.41	(1)													P
<b>TOTAL NATIVE PERENNIAL GRASSES (c)</b>			8.2 (8.3)	100.0	25.4 (25.3)	9.3	25.7	12(2)	6(1)	(1)	---	10(1)	1	---	7(1)	---	---	9	5	---	---
<b>INTRODUCED PERENNIAL GRASSES (cool)</b>																					
<i>Ceratochloa catineta</i>	CERATOCHELOA MARGINATA, BROMUS MARGINATUS, B. POLYANTHUS	MOUNTAIN BROME	0.00	50.00	0.00	0.00	0.00		P						P				P		
<i>Dactylis glomerata</i>		ORCHARD GRASS	0.17	16.67	0.52	0.17	0.41														1
<i>Poa compressa</i>		CANADA BLUEGRASS	0.00	33.33	0.00	0.00	0.00	P													
<b>TOTAL INTRO. PERENNIAL GRASSES (c)</b>			0.2	66.7	0.5	0.2	0.5	P	P	---	---	---	---	---	P	---	---	1	---	---	---
<b>NATIVE PERENNIAL GRASSES (warm)</b>																					
<i>Chondrosium gracile</i>	BOULELOUA GRACILIS	BLUE GRAMA GRASS	0.33	66.67	1.04	0.33	0.83	P	P			P									2
<i>Muhlenbergia montana</i>		MOUNTAIN MUHLY	0.00	16.67	0.00	0.00	0.00					P									
<b>TOTAL NATIVE PERENNIAL GRASSES (w)</b>			0.3	66.7	1.0	0.3	0.9	P	P	---	---	P	---	---	---	---	---	2	---	---	---
<b>NATIVE SHRUBS</b>																					
<i>Arotostaphylos uva-ursi</i>		KINKINNICK	0.17	50.00	0.52	0.33	0.83	1(1)							P				P		
<i>Ceanothus fendleri</i>		BUCKBRUSH	5.00 (5.17)	100.00	15.54 (15.66)	5.67	14.11	3	4			8			1	1		11	3(2)	(1)	
<i>Physocarpus monogynus</i>		NINEBARK	0.00	16.67	0.00	0.00	0.00								P						
<i>Ribes cereum</i>		WAX CURRANT	0.33	100.00	1.04	0.33	0.83	P	2			P			P				P	P	
<i>Rosa arkansana</i>		ARKANSAS ROSE	0.17	16.67	0.52	0.17	0.41								1						
<i>Rosa woodsii</i>		WOOD'S ROSE	0.00	16.67	0.00	0.00	0.00								P						
<i>Rubus idaeus ssp. melanoclasius</i>		AMERICAN RED RASPBERRY	0.33	66.67	1.04	0.33	0.83	1	1						P						P
<b>TOTAL NATIVE SHRUBS</b>			6.0 (6.2)	100.0	18.7 (18.7)	6.8	18.6	5(1)	7	---	---	8	---	---	2	1	---	11	3(2)	(1)	---
<b>NATIVE TREES</b>																					
<i>Populus tremuloides</i>	POPULUS TREMULA	QUAKING ASPEN	0.00	16.67	0.00	0.00	0.00								P						
<i>Pseudotsuga menziesii deac</i>		DOUGLAS FIR	3.50 (1.17)	83.33	10.88 (3.54)	3.83	9.54	[4]	[1]		[1]		[1]	[6]		[1]	[6]				[3]
<b>TOTAL NATIVE TREES</b>			0.0	16.7	0.0	0.0	0.0	---	---	---	---	---	---	---	P	---	---	---	---	---	---
Standing dead		STANDING DEAD	6.17 (0.00)	100.00		6.50		5	1			5			13	1		3	10	1	
Litter		LITTER	35.50 (36.50)	100.00		36.67		59	28			38	4		27	1		36	25(1)	1	
Bare soil		BARE SOIL	20.00 (20.17)	100.00		20.17		4	31			18			21	1		21	25		
Rock		ROCK	2.67	66.67		2.67			7			2			4				3		
<b>TOTALS</b>			96.5 (93.5)			106.2		96	98			94			94	5		100	97		
<b>TOTALS (LAYER)</b>								96	98	1	0	94	5	0	94	5	0	100	97	3	0
<b>TOTAL VEGETATION COVER (LAYER)</b>								28(4)	31(1)	1(1)	0(1)	31(2)	1	0	29(3)	2	0	40(1)	34(6)	1(1)	0
<b>TOTAL VEGETATION COVER</b>			32.2 (34.2)		100.0 (103.5)	40.2 (s=14.7)	100.0	28(4)	31(4)			31(3)			29(5)			40(1)	34(6)		
<b>GROUND COVER (Litter+Rock+Veg+St.Dead)</b>			Std.Dev.= 4.4 76.5 (78.7)			86.0 (83.2)		92(4)	67(1)	1(1)	0(1)	76(2)	5	0	73(3)	4	0	79(1)	72(7)	3(1)	0
<b>SPECIES DENSITY (# of species/100 sq.m.)</b>								45	38			32			37			40	37		
<b>(AVERAGE= 38.2 Std.Dev.= 4.3)</b>																					
<b>SPECIES DENSITY (LAYER)</b>								45	38	2	1	32	1	0	37	2	0	40	37	2	0

(#) = second hit  
 (#) = cover value if tree canopy is excluded  
 (#) = hit on standing dead tree species

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	02D	04D	05D	05DWU	05U	07D	11D	11DWU	11U
<b>NATIVE ANNUAL &amp; BIENNIAL FORBS</b>																
<i>Bahia dissecta</i>		CUTLEAF	0.40	40.00	0.85	0.40	0.76		2	P						
<i>Gnndelia squarrosa</i>		GUMWEED	0.00	80.00	0.00	0.00	0.00		P	P			P	P		
<i>Machaeranthera bigelovii</i>		BIGELOW ASTER	0.00	20.00	0.00	0.00	0.00			P						
<i>Pterogonum alatum</i>	ERIOGONUM ALATUM	WINGED BUCKWHEAT	0.00	40.00	0.00	0.00	0.00	P						P		
<i>Silene antirrhina</i>		SLEEPY CATCHFLY	0.00	20.00	0.00	0.00	0.00	P								
<b>TOTAL NATIVE ANN. &amp; BIEN. FORBS</b>			0.4	100.0	0.8	0.4	0.8	P	2	P	---	---	P	P	---	---
<b>INTRODUCED ANNUAL &amp; BIENNIAL FORBS</b>																
<i>Acosta diffusa</i>	CENTAUREA DIFFUSA	TUMBLE KNAPWEED	1.80 (2.00)	20.00	3.81 (4.20)	2.00	3.80							9	1	
<i>Alyssum minus</i>		ALYSSUM	0.80	80.00	1.69	0.80	1.52	P	1				P	3		
<i>Camelina microcarpa</i>		LITTLEPOD FALSEFLAX	0.00	60.00	0.00	0.00	0.00	P					P	P		
<i>Lactuca serriola</i>		PRICKLY LETTUCE	0.00	40.00	0.00	0.00	0.00	P						P		
<i>Lappula redowskii</i>		EARLY STICKSEED	0.00	20.00	0.00	0.00	0.00	P								
<i>Sisymbrium altissimum</i>		JIM HILL MUSTARD	0.00	40.00	0.00	0.00	0.00	P						P		
<i>Tragopogon dubius ssp. major</i>		YELLOW SALSIFY	0.00	80.00	0.00	0.00	0.00	P	P				P	P		
<i>Verbascum thapsus</i>		MULLEIN	1.20	80.00	2.54	1.20	2.28		4	1			1	P		
<b>TOTAL INTRO. ANN. &amp; BIEN. FORBS</b>			3.8 (4.0)	100.0	8.1 (8.4)	4.0	7.7	P	5	1	---	---	1	12	1	---
<b>INTRODUCED ANNUAL GRASSES</b>																
<i>Anisantha tectorum</i>	BROMUS TECTORUM	CHEAT GRASS	17.40	100.00	36.86	18.20	34.60	19	22(1)	20			13(1)	13(2)		
<i>Bromus japonicus</i>		JAPANESE BROME	0.00	20.00	0.00	0.00	0.00							P		
<i>Cylindropyrum cylindricum</i>	AEGILOPS CYLINDRICA	GOAT GRASS	0.00	20.00	0.00	0.00	0.00	P								
<b>TOTAL INTRO. ANN. GRASSES</b>			17.4	100.0	36.9	18.2	35.0	19	22(1)	20	---	---	13(1)	13(2)	---	---
<b>NATIVE PERENNIAL FORBS</b>																
<i>Achillea lanulosa</i>		WESTERN YARROW	0.20	60.00	0.42	0.20	0.38		P	P				1		
<i>Apocynum androsaemifolium</i>		SPREADING DOGBANE	0.00	20.00	0.00	0.00	0.00			P						
<i>Artemisia frigida</i>		FRINGED SAGE	6.00 (6.20)	80.00	12.71 (13.03)	7.00	13.31	11		14(1)	1		5(3)	P		
<i>Artemisia ludoviciana</i>		PASTURE SAGE	1.00	80.00	2.12	1.80	3.42		1	3(4)			1	P		
<i>Aster porteri</i>		PORTER'S ASTER	0.00	80.00	0.00	0.00	0.00	P	P	P			P			
<i>Astragalus laxmannii</i>	ASTRAGALUS ADSURGENS VAR. ROBUSTIER	LAXMANN'S MILKVETCH	0.00	20.00	0.00	0.00	0.00						P			
<i>Campanula rotundifolia</i>		HAREBELL	0.00	20.00	0.00	0.00	0.00			P						
<i>Cirsium ochrocentrum</i>	CIRSIUM MEGACEPHALUM	THISTLE	0.00	40.00	0.00	0.00	0.00	P						P		
<i>Drymocallis fissa</i>	POTENTILLA FISSA	BIGFLOWER CINQUEFOIL	0.00	60.00	0.00	0.00	0.00	P	P					P		
<i>Eriogonum umbellatum var. umbellatum</i>		WILD BUCKWHEAT	0.80	60.00	1.69	0.80	1.52		4	P				P		
<i>Erysimum capitatum</i>		COAST WALLFLOWER	0.00	20.00	0.00	0.00	0.00		P							
<i>Gaillardia aristata</i>		BLANKET FLOWER	0.00	80.00	0.00	0.00	0.00	P		P			P	P		
<i>Geranium caespitosum ssp. caespitosum</i>		SMALL-LEAF WILD GERANIUM	0.80	80.00	1.69	1.00	1.90	3	(1)				1	P		
<i>Harbouria trachypleura</i>		WHISKBROOM PARSLEY	0.00	40.00	0.00	0.00	0.00			P			P			
<i>Helianthus pumilus</i>		SUNFLOWER	0.20	60.00	0.42	0.40	0.76	(1)					P	1		
<i>Heterotheca foliosa</i>		GOLDENASTER	5.40	80.00	11.44	5.60	10.65	7		P			11	9(1)		
<i>Heterotheca villosa</i>	HETEROTHECA HORRIDA, CHRYSOPSIS VILLOSA	HARRY GOLDEN ASTER	0.40	40.00	0.85	0.40	0.76		2	P						
<i>Mertensia lanceolata</i>	MERTENSIA VIRIDIS, M. BAKERI	LANCELEAF BLUEBELLS	0.00	20.00	0.00	0.00	0.00							P		
<i>Penstemon virens</i>		GREEN BEARD-TONGUE	0.60	80.00	1.27	0.60	1.14	P		P			2	1		
<i>Phacelia heterophylla</i>		VARIABLE SCORPIONWEED	0.00	40.00	0.00	0.00	0.00		P	P						
<i>Pulsatilla ludoviciana</i>	P. PATENS SSP. MULTIFIDA	PASQUEFLOWER	0.00	20.00	0.00	0.00	0.00			P						
<i>Rumex triangulivalvis</i>	RUMEX SALICIFOLIUS	WILLOW DOCK	0.00	20.00	0.00	0.00	0.00							P		
<i>Solidago simplex var. simplex</i>	SOLIDAGO SPATHULATA VAR. NEOMEXICANA	MT. ALBERT GOLDENROD	1.20	40.00	2.54	1.40	2.66		1	5(1)						
<b>TOTAL NATIVE PERENNIAL FORBS</b>			16.6 (16.8)	100.0	35.2 (35.3)	19.2	36.9	21(1)	8(1)	22(6)	1	---	20(3)	12(1)	---	---
<b>INTRODUCED PERENNIAL FORBS</b>																
<i>Breca arvensis</i>	CIRSIUM ARVENSE	CANADA THISTLE	0.00	20.00	0.00	0.00	0.00	P								
<b>TOTAL INTRO. PERENNIAL FORBS</b>			0.0	20.0	0.0	0.0	0.0	P	---	---	---	---	---	---	---	---
<b>NATIVE PERENNIAL GRASSES (cool)</b>																
<i>Bromopsis lanatipes</i>	BROMUS LANATIPES	WOOLY BROME	0.00	20.00	0.00	0.00	0.00			P						
<i>Carex pensylvanica ssp. heliophila</i>	CAREX HELIOPHILA	SUN SEDGE	1.60	100.00	3.39	2.40	4.56	2(1)	2	P			1(1)	3(2)		
<i>Elymus elymoides</i>	SITANION HYSTRIX	BOTTLEBRUSH SQUIRRELTAIL	0.20	80.00	0.42	0.20	0.38		P	P			P	1		
<i>Elymus lanceolatus fm. albicans</i>	AGROPYRON DASYSTACHYUM A. ALBICANS, AGROPYRON RIPARIUM	MONTANA WHEATGRASS	0.80	40.00	1.69	0.80	1.52			P			4			
<i>Elymus trachycaulus</i>	AGROPYRON TRACHYCAULUM	SLENDER WHEATGRASS	1.00	20.00	2.12	1.00	1.90							5		
<i>Hesperostipa comata</i>	STIPA COMATA	NEEDLE-AND-THREAD GRASS	0.60	80.00	1.27	0.60	1.14	1	1				P	1		
<i>Koeleria macrantha</i>	KOELERIA CRISTATA K. PYRAMIDATA K. GRACILIS	JUNE GRASS	0.00	60.00	0.00	0.00	0.00	P	P				P			
<i>Leucopoa kingii</i>		SPIKE FESCUE	0.00	60.00	0.00	0.00	0.00			P			P	P		
<i>Poa agassizensis</i>		AGASSIZ BLUEGRASS	0.00	40.00	0.00	0.00	0.00	P		P						
<b>TOTAL NATIVE PERENNIAL GRASSES (c)</b>			4.2	100.0	8.9	5.0	9.6	3(1)	3	P	---	---	5(1)	10(2)	---	---

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	02D	04D	05D	05DWU	05U	07D	11D	11DWU	11U
<b>INTRODUCED PERENNIAL GRASSES (cool)</b>																
<i>Ceratochloa carinata</i>	CERATOCHLOA MARGINATA, BROMUS MARGINATUS, B. POLYANTHUS	MOUNTAIN BROME	0.00	20.00	0.00	0.00	0.00							P		
<i>Poa compressa</i>		CANADA BLUEGRASS	0.00	20.00	0.00	0.00	0.00			P						
<b>TOTAL INTRO. PERENNIAL GRASSES (c)</b>																
			0.0	40.0	0.0	0.0	0.0	---	---	P	---	---	---	P	---	---
<b>NATIVE PERENNIAL GRASSES (warm)</b>																
<i>Chondrosium gracile</i>	BOUTELOUA GRACILIS	BLUE GRAMA GRASS	0.00	60.00	0.00	0.20	0.38	P	(1)				P			
<i>Muhlenbergia montana</i>		MOUNTAIN MUHLY	0.80	60.00	1.69	0.80	1.52	P	1				3			
<b>TOTAL NATIVE PERENNIAL GRASSES (w)</b>																
			0.8	60.0	1.7	1.0	1.9	P	1(1)	---	---	---	3	---	---	---
<b>NATIVE SHRUBS</b>																
<i>Ceanothus fendleri</i>		BUCKBRUSH	2.80	100.00	5.93	2.80	5.32	2	6	P			4	2		
<i>Cercocarpus montanus</i>		BIRCHLEAF MOUNTAIN MAHOGANY	0.60	60.00	1.27	0.80	1.52		2	P		(1)		1		
<i>Physocarpus monogynus</i>		NINEBARK	0.20	20.00	0.42	0.20	0.38		1							
<i>Ribes cereum</i>		WAX CURRANT	0.00	60.00	0.00	0.00	0.00	P	P					P		
<i>Rosa arkansana</i>		ARKANSAS ROSE	0.20	60.00	0.42	0.20	0.38	P		P						
<i>Rubus idaeus ssp. melanolasius</i>		AMERICAN RED RASPBERRY	0.20	40.00	0.42	0.20	0.38	1		P				1		
<b>TOTAL NATIVE SHRUBS</b>																
			4.0	100.0	8.5	4.2	8.1	3	9	P	---	(1)	4	4	---	---
<b>NATIVE TREES</b>																
<i>Pinus ponderosa ssp. scopulorum (dead)</i>		PONDEROSA PINE (DEAD)	0.40 (0.20)	20.00	0.85 (0.42)	0.60	1.14					[1]	[2]			
<b>TOTAL NATIVE TREES</b>																
			0.0		0.0	0.0	0.0									
Standing dead		STANDING DEAD	6.00 (0.00)	80.00		6.00		1		9			9	10		1
Litter		LITTER	27.20	100.00		27.20		25	33	29			23	26		
Bare soil		BARE SOIL	17.80	100.00		17.80		28	15	13			21	12		
Rock		ROCK	1.40	60.00		1.40			2	4			1			
<b>TOTALS</b>																
			99.6 (94.2)			105.0		100	100	98			100	100		
<b>TOTALS (LAYER)</b>																
<b>TOTAL VEGETATION COVER (LAYER)</b>																
<b>TOTAL VEGETATION COVER</b>																
			47.2 (47.8)		100.0 (100.4)	52.6 (s=22.8)	100.0	46(2)	50(3)	43(6)	1	0(1)	46(5)	51(5)	1	0
			Std.Dev = 3.3					46(2)	50(3)	43(6)			46(5)	51(6)		
<b>GROUND COVER (Litter+Rock+Veg+St. Dead)</b>																
			81.8 (82.0)			87.2 (86.4)		72(2)	85(3)	85(6)	1	0(1)	79(5)	87(5)	1	1
<b>SPECIES DENSITY (# of species/100 sq.m.)</b>																
<b>(AVERAGE= 29.2 Std.Dev = 4.0)</b>																
<b>SPECIES DENSITY (LAYER)</b>																
								31	25	31			25	34		
								31	25	31	1	1	25	34	1	0

(#) = second hit  
 (#) = cover value if tree canopy is excluded  
 (#) = hit on standing dead tree species

## DATA FROM FILE Group D

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	03D	15D	15DWU	15U	17D	17DWU	17U
<b>NATIVE ANNUAL &amp; BIENNIAL FORBS</b>														
<i>Aclroslia dispersa</i>	MENTZELIA	BUSHY BLAZING STAR	0	33.33	0	0	0		P					
<i>Chenopodium leptophyllum</i>		NARROWLEAF GOOSEFOOT	0	66.67	0	0	0	P	P					
<i>Collinsia parviflora</i>		BABY BLUE-EYES	0	66.67	0	0	0		P				P	
<i>Epilobium brachycarpum</i>	EPILOBIUM PANICULATUM	BIGFRUIT WILLOWHERB	0	33.33	0	0	0		P					
<i>Erigeron divergens</i>		SPREADING FLEABANE	0	33.33	0	0	0		P					
<i>Grindelia squarrosa</i>		GUMWEED	0	33.33	0	0	0							P
<i>Machaeranthera bigelovii</i>		BIGELOW ASTER	0	66.67	0	0	0		P					P
<i>Pterogonum alatum</i>	ERIOGONUM ALATUM	WINGED BUCKWHEAT	0	66.67	0	0	0	P						P
<b>TOTAL NATIVE ANN. &amp; BIEN. FORBS</b>			0	100	0	0	0	P	P	---	---	P	---	---
<b>INTRODUCED ANNUAL &amp; BIENNIAL FORBS</b>														
<i>Alyssum minus</i>		ALYSSUM	0	67	1	0	1	1						P
<i>Coryza canadensis</i>		HORSEWEED	0	33.33	0	0	0		P					
<i>Descurainia sophia</i>		FLUXWEED TANSYMUSTARD	0	33.33	0	0	0		P					
<i>Lactuca serriola</i>		PRICKLY LETTUCE	0	66.67	0	0	0	P	P					
<i>Sisymbrium altissimum</i>		JIM HILL MUSTARD	0.67	33.33	1.22	0.67	1.06	2						
<i>Tragopogon dubius ssp. major</i>		YELLOW SALSIFY	0	66.67	0	0	0	P	P					
<i>Turnitis glabra</i>	ARABIS GLABRA	TOWER MUSTARD	0	33.33	0	0	0							P
<i>Verbascum thapsus</i>		MULLEIN	0.67	100	1.22	0.67	1.06	1	1					P
<b>TOTAL INTRO. ANN. &amp; BIEN. FORBS</b>			2	100	3	2	3	4	1	---	---	P	---	---
<b>INTRODUCED ANNUAL GRASSES</b>														
<i>Anisantha tectorum</i>	BROMUS TECTORUM	CHEATGRASS	22.00 (22.33)	100	40.24 (40.61)	24	38	25(2)	31(2)	1			10(1)	
<b>TOTAL INTRO. ANN. GRASSES</b>			22.0 (22.3)	100	40.2 (40.6)	24	39	25(2)	31(2)	1	---	---	10(1)	---
<b>NATIVE PERENNIAL FORBS</b>														
<i>Achillea lanulosa</i>		WESTERN YARROW	0	67	0	0	0	P	P					
<i>Aletes acaulis</i>		STEMLESS INDIAN PARSLEY	0	33.33	0	0	0							P
<i>Allium cernuum</i>		NODDING ONION	0	66.67	0	0	0	P						P
<i>Amerosedum lanceolatum</i>	SEDUM LANCEOLATUM	YELLOW STONECROP	0	33.33	0	0	0							P
<i>Artemisia frigida</i>		FRINGED SAGE	1.33	100	2.44 (2.42)	1.67	2.66	4	P					(1)
<i>Artemisia ludoviciana</i>		PASTURE SAGE	0.33	100	0.61	1	1.6	P	(1)				1(1)	(1)
<i>Aster porteri</i>		PORTER'S ASTER	0.33	100	0.61	0.33	0.53	P	P				1	
<i>Campanula rotundifolia</i>		HAREBELL	0	66.67	0	0	0	P	P					
<i>Cerastium strictum</i>	CERASTIUM ARVENSE	MOUSE-EAR	1	33.33	1.83	1	1.6		3					
<i>Erigeron compositus</i>		FLEABANE	0	33.33	0	0	0							P
<i>Eriogonum umbellatum var. umbellatum</i>		WLD BUCKWHEAT	1	33.33	1.83	1	1.6							3
<i>Erysimum capitatum</i>		COAST WALLFLOWER	0	66.67	0	0	0		P					P
<i>Gaillardia aristata</i>		BLANKETFLOWER	0	33.33	0	0	0							P
<i>Gastrolychnis drummondii</i>	MELANDRUM DRUMMONDII, LYCHNIS DRUMMONDII	DRUMMOND CAMPION	0	33.33	0	0	0							P
<i>Geranium caespitosum ssp. caespitosum</i>		SMALL-LEAF WILD GERANIUM	0.33	66.67	0.61	0.33	0.53	P						1
<i>Harboursia trachyleura</i>		WHISKBROOM PARSLEY	0	33.33	0	0	0							P
<i>Helianthus pumilus</i>		SUNFLOWER	0.67	100	1.22	0.67	1.06	1	1					P
<i>Heterotheca foliosa</i>		GOLDENASTER	3.33	100	6.1	3.67	5.85	9(1)	1					P
<i>Lesquerella montana</i>		BLADDERPOD	0	33.33	0	0	0							P
<i>Microseris nutans</i>		MICROSERIS	0.33	33.33	0.61	0.33	0.53		1					
<i>Packera fendleri</i>	SENECIO FENDLERI	FENDLER'S RAGWORT	0	33.33	0	0	0							P
<i>Paronychia jamesii</i>		NAILWORT	0	33.33	0	0	0							P
<i>Pensemon virens</i>		GREEN BEARD-TONGUE	0.67 (1.00)	66.67	1.22 (1.82)	1	1.6		P				2	1
<i>Phacelia heterophylla</i>		VARIABLE SCORPIONWEED	0	33.33	0	0	0	P						
<i>Scutellaria brittonii</i>		SKULLCAP	0	33.33	0	0	0							P
<b>TOTAL NATIVE PERENNIAL FORBS</b>			9.3 (9.7)	100	17.1 (17.6)	11	18	14(1)	6(1)	---	---	---	8(1)	1(1) ---
<b>INTRODUCED PERENNIAL FORBS</b>														
<i>Breera arvensis</i>	CIRSIUM ARVENSE	CANADA THISTLE	0	33	0	0	0	P						
<b>TOTAL INTRO. PERENNIAL FORBS</b>			0	33	0	0	0	P	---	---	---	---	---	---
<b>NATIVE PERENNIAL GRASSES (cool)</b>														
<i>Bromopsis lanatipes</i>	BROMUS LANATIPES	WOOLY BROME	0	67	0	0	0	P						P
<i>Carex pensylvanica ssp. heliophila</i>	CAREX HELIOPHILA	SUN SEDGE	1.67	66.67	3.05	2	3.19	2	3(1)					
<i>Elymus elymoides</i>	SITANION HYSTRIX	BOTTLEBRUSH SQUIRRELTAIL	2.67	66.67	4.88	3	4.79	8(1)						P
<i>Elymus lanceolatus fm. albicans</i>	AGROPYRON DASYSTACHYUM, A. ALBICANS, AGROPYRON RIPARIUM	MONTANA WHEATGRASS	0.67	66.67	1.22	0.67	1.06	2						P
<i>Festuca arizonica</i>		ARIZONA FESCUE	0.33	33.33	0.61	0.33	0.53		1					
<i>Hesperostipa comata</i>	STIPA COMATA	NEEDLE-AND-THREAD GRASS	0	33.33	0	0	0	P						
<i>Koeleria macrantha</i>	KOELERIA CRISTATA, K. PYRAMIDATA, K. GRACILIS	JUNEGRASS	0.33	33.33	0.61	0.33	0.53	1						
<i>Leucopoa kingii</i>		SPIKE FESCUE	0.67	100	1.22	0.67	1.06	1	1					P
<b>TOTAL NATIVE PERENNIAL GRASSES (c)</b>			6	100	12	7	12	14(1)	5(1)	---	---	---	P	---

DATA FROM FILE Group D

SCIENTIFIC NAME	SYNONYM	COMMON	AVERAGE COVER (%)	FREQUENCY (%)	RELATIVE VEGETATION COVER (%)	AVERAGE COVER-ALL (%)	RELATIVE VEGETATION COVER-ALL (%)	03D	15D	15DWU	15U	17D	17DWU	17U
<b>NATIVE PERENNIAL GRASSES (warm)</b>														
<i>Muhlenbergia montana</i>		MOUNTAIN MUHLY	1	67	1	1	1		1			1		
<b>TOTAL NATIVE PERENNIAL GRASSES (w)</b>			1	67	1	1	1	---	1	---	---	1	---	---
<b>NATIVE SHRUBS</b>														
<i>Ceanothus fendleri</i>		BUCKBRUSH	8.00 {8.33}	100	14.63 {15.15}	10	15	3(1)	14(2)			7(1)	1	
<i>Cercocarpus montanus</i>		BIRCHLE AF MOUNTAIN MAHOGANY	5.67	66.67	10.37	6	9.57		4			13(1)		
<i>Padus virginiana ssp. melanocarpa</i>	PRUNUS VIRGINIANA SSP. MELANOCARPA	CHOKECHERRY	0	33.33	0	0	0		P					
<i>Physocarpus monogynus</i>		NINEBARK	0	33.33	0	0	0		P					
<i>Ribes cereum</i>		WAX CURRANT	0.33	100	0.61	0.33	0.53	P	P			1		
<i>Yucca glauca</i>		SP ANISH BAYONET	0	33.33	0	0	0					P		
<b>TOTAL NATIVE SHRUBS</b>			14.0 {14.3}	100	25.6 {26.1}	16	26	3(1)	18(2)	---	---	21(2)	1	---
<b>NATIVE TREES</b>														
<i>Pinus ponderosa ssp. scopulorum</i>		PONDEROSA PINE	0.33 {0.00}	33	1	0	1							1
<i>Pseudotsuga menziesii</i>		DOUGLAS-FIR	0.33 {0.00}	33.33	0.61	0.33	0.53							1
<i>Pseudotsuga menziesii dead</i>		DOUGLAS FIR	0.67 {0.00}	33.33		0.67	1.06							[2]
<i>Sabina scopulorum dead</i>	JUNIPERUS SCOPULORUM	ROCKY MOUNTAIN JUNIPER	1.00 {0.67}	33.33		1	1.6		[2]		[1]			
<b>TOTAL NATIVE TREES</b>			0.7 {0.0}	33	1	1	1	---	---	---	---	---	---	2
Standing dead		STANDING DEAD	1.67 {0.00}	66.67		1.67			2			3		
Litter		LITTER	22.00 {22.33}	100		22.33		29	17			20	1	
Bare soil		BARE SOIL	14.33 {14.67}	100		14.67		8	10			25	1	
Rock		ROCK	5.67	100		5.67		3	6			8		
<b>TOTALS</b>			98.3 {98.3}			107		100	97			98		
<b>TOTALS (LAYER)</b>														
<b>TOTAL VEGETATION COVER (LAYER)</b>								100	97	1	0	96	4	2
<b>TOTAL VEGETATION COVER</b>			54.7 {55.7}		100.0 {101.2}	62.7 (s=37.5)	100	80(5)	82(6)	1	0	40(4)	2(1)	2
<b>GROUND COVER (Litter+Rock+Veg+St. Dead)</b>			84.0 {84.7}			92.3 {90.7}		80(5)	82(7)			42(7)		
<b>SPECIES DENSITY (# of species/100 sq.m.)</b>								28	32			39		
<b>(AVERAGE= 33.0 Std.Dev.= 5.6)</b>														
<b>SPECIES DENSITY (LAYER)</b>								28	32	1	0	36	3	2

(#) = second hit  
 {#} = cover value if tree canopy is excluded  
 [#] = hit on standing dead tree species

Appendix 2. Species Importance in 2002 and 2007 based on 18 samples.

Scientific Name	Relative Importance (%)	Constancy (%)	Average Cover All Hits (%)
<i>Pinus ponderosa</i> ssp. <i>scopulorum</i> (dead)	100.000	66.67	5.61
*** <i>Carex pensylvanica</i> ssp. <i>heliophila</i>	65.923	77.78	3.17
<i>Ceanothus fendleri</i>	53.278	94.44	2.11
<i>Phacelia heterophylla</i>	53.237	77.78	2.56
<i>Pseudotsuga menziesii</i> (dead)	42.931	55.56	2.89
<i>Verbascum thapsus</i>	36.360	94.44	1.44
<i>Sisymbrium altissimum</i>	28.103	61.11	1.72
<i>Chenopodium simplex</i>	27.658	77.78	1.33
*** <i>Anisantha tectorum</i>	24.716	72.22	1.28
<i>Elymus trachycaulus</i>	19.338	33.33	2.17
**** <i>Ceratochloa carinata</i>	18.357	33.33	2.06
<i>Penstemon virens</i>	15.923	88.89	0.67
<i>Artemisia ludoviciana</i>	12.937	72.22	0.67
<i>Geranium caespitosum</i> ssp. <i>caespitosum</i>	12.685	77.78	0.61
<i>Carex</i> spp.	11.096	50.00	0.83
<i>Corydalis aurea</i>	9.953	55.56	0.67
<i>Artemisia frigida</i>	8.957	50.00	0.67
<i>Campanula rotundifolia</i>	8.110	77.78	0.39
<i>Chenopodium leptophyllum</i>	8.110	77.78	0.39
<i>Grindelia squarrosa</i>	7.248	44.44	0.61
<i>Triticum aestivum</i> x <i>Elytrigia elongata</i>	7.189	61.11	0.44
<i>Helianthus pumilus</i>	6.952	66.67	0.39
<i>Breea arvensis</i>	6.343	38.89	0.61
<i>Apocynum androsaemifolium</i>	6.297	22.22	1.06
<i>Chondrosium gracile</i>	5.941	44.44	0.50
<i>Epilobium brachycarpum</i>	5.882	66.67	0.33
<i>Leucopoa kingii</i>	5.882	66.67	0.33
<i>Astragalus miser</i> var. <i>oblongifolius</i>	5.407	72.22	0.28
*** <i>Populus tremuloides</i>	4.931	22.22	0.83
<i>Pseudotsuga menziesii</i>	4.876	5.56	3.28
<i>Physocarpus monogynus</i>	4.634	44.44	0.39
<i>Heterotheca villosa</i>	4.412	50.00	0.33
<i>Lactuca serriola</i>	3.922	66.67	0.22
<i>Solidago simplex</i> var. <i>simplex</i>	3.743	50.00	0.28
<i>Elymus lanceolatus</i> fm. <i>albicans</i>	3.431	38.89	0.33
<i>Sabina scopulorum</i> (dead)	3.327	22.22	0.56
<i>Alyssum minus</i>	3.268	27.78	0.44
<i>Erysimum capitatum</i>	3.030	66.67	0.17
<i>Elymus elymoides</i>	2.614	44.44	0.22

<i>Rubus idaeus</i> ssp. <i>melanolasius</i>	2.525	55.56	0.17
<i>Silene antirrhina</i>	2.525	55.56	0.17
<i>Gaillardia aristata</i>	2.273	50.00	0.17
<i>Heterotheca foliosa</i>	2.080	27.78	0.28
<i>Drymocallis fissa</i>	1.961	66.67	0.11
<i>Ribes cereum</i>	1.961	66.67	0.11
<i>Aster porteri</i>	1.797	61.11	0.11
<i>Cercocarpus montanus</i>	1.663	22.22	0.28
<i>Rosa arkansana</i>	1.515	33.33	0.17
<i>Aletes acaulis</i>	1.144	38.89	0.11
<i>Astragalus laxmannii</i>	1.144	38.89	0.11
<i>Carduus nutans</i> ssp. <i>macrolepis</i>	1.144	38.89	0.11
<i>Oenothera caespitosa</i> ssp. <i>caespitosa</i>	1.144	38.89	0.11
<i>Dracocephalum parviflorum</i>	0.981	16.67	0.22
<i>Achillea lanulosa</i>	0.980	61.11	0.06
<i>Acrolasia dispersa</i>	0.980	33.33	0.11
<i>Pulsatilla ludoviciana</i>	0.802	50.00	0.06
<i>Scutellaria brittonii</i>	0.802	50.00	0.06
<i>Helianthus annuus</i>	0.653	22.22	0.11
*** <i>Androsace septentrionalis</i>	0.624	38.89	0.06
<i>Taraxacum officinale</i>	0.624	38.89	0.06
<i>Harbouria trachypleura</i>	0.535	33.33	0.06
<i>Muhlenbergia montana</i>	0.505	11.11	0.17
<i>Cirsium ochrocentrum</i>	0.490	16.67	0.11
<i>Bassia sieversiana</i>	0.446	27.78	0.06
<i>Poa agassizensis</i>	0.446	27.78	0.06
<i>Hesperostipa comata</i>	0.356	22.22	0.06
<i>Poa compressa</i>	0.327	11.11	0.11
<i>Populus tremuloides</i> (dead)	0.327	11.11	0.11
<i>Arctostaphylos uva-ursi</i>	0.267	16.67	0.06
<i>Lappula redowskii</i>	0.267	16.67	0.06
<i>Liatis punctata</i>	0.267	16.67	0.06
<i>Symphoricarpos rotundifolius</i>	0.267	16.67	0.06
<i>Astragalus agrestis</i>	0.178	11.11	0.06
<i>Monarda fistulosa</i> var. <i>menthifolia</i>	0.178	11.11	0.06
<i>Oxytropis lambertii</i>	0.178	11.11	0.06
<i>Verbena bracteata</i>	0.178	11.11	0.06
Moss	0.164	5.56	0.11
<i>Mertensia lanceolata</i>	0.134	50.00	0.01
<i>Eriogonum umbellatum</i> var. <i>umbellatum</i>	0.104	38.89	0.01
<i>Cirsium vulgare</i>	0.089	5.56	0.06

<i>Festuca brachyphylla</i> ssp. <i>coloradensis</i>	0.089	5.56	0.06
<i>Machaeranthera bigelovii</i>	0.089	5.56	0.06
<i>Koeleria macrantha</i>	0.089	33.33	0.01
<i>Polygonum douglasii</i>	0.074	27.78	0.01
<i>Oreobatus deliciosus</i>	0.059	22.22	0.01
<i>Allium cernuum</i>	0.045	16.67	0.01
<i>Astragalus shortianus</i>	0.045	16.67	0.01
<i>Conyza canadensis</i>	0.045	16.67	0.01
<i>Galium septentrionale</i>	0.045	16.67	0.01
*** <i>Lesquerella montana</i>	0.045	16.67	0.01
<i>Nicotiana attenuata</i>	0.045	16.67	0.01
<i>Packera fendleri</i>	0.045	16.67	0.01
<i>Penstemon glaber</i>	0.045	16.67	0.01
<i>Potentilla hippiana</i>	0.045	16.67	0.01
<i>Tragopogon dubius</i> ssp. <i>major</i>	0.045	16.67	0.01
<i>Acer glabrum</i>	0.030	11.11	0.01
<i>Acosta diffusa</i>	0.030	11.11	0.01
<i>Amerosedum lanceolatum</i>	0.030	11.11	0.01
<i>Arnica fulgens</i>	0.030	11.11	0.01
<i>Astragalus</i> spp.	0.030	11.11	0.01
<i>Bromopsis lanatipes</i>	0.030	11.11	0.01
<i>Bromus japonicus</i>	0.030	11.11	0.01
<i>Camelina microcarpa</i>	0.030	11.11	0.01
<i>Cerastium strictum</i>	0.030	11.11	0.01
<i>Collomia linearis</i>	0.030	11.11	0.01
<i>Cylindropyrum cylindricum</i>	0.030	11.11	0.01
<i>Euphorbia</i> spp.	0.030	11.11	0.01
<i>Frasera speciosa</i>	0.030	11.11	0.01
<i>Gaura mollis</i>	0.030	11.11	0.01
<i>Lupinus argenteus</i>	0.030	11.11	0.01
<i>Physaria vitulifera</i>	0.030	11.11	0.01
<i>Rosa woodsii</i>	0.030	11.11	0.01
<i>Sporobolus cryptandrus</i>	0.030	11.11	0.01
<i>Achnatherum nelsonii</i>	0.015	5.56	0.01
<i>Agrostis scabra</i>	0.015	5.56	0.01
<i>Antennaria rosea</i>	0.015	5.56	0.01
<i>Arabis hirsuta</i>	0.015	5.56	0.01
<i>Asclepias stenophylla</i>	0.015	5.56	0.01
<i>Asclepias viridiflora</i>	0.015	5.56	0.01
<i>Astragalus drummondii</i>	0.015	5.56	0.01
<i>Bahia dissecta</i>	0.015	5.56	0.01
<i>Boechea fendleri</i>	0.015	5.56	0.01
<i>Carex</i> sp. 1	0.015	5.56	0.01

<i>Chenopodium foliosum</i>	0.015	5.56	0.01
<i>Chrysothamnus parryi</i>	0.015	5.56	0.01
<i>Collinsia parviflora</i>	0.015	5.56	0.01
<i>Cystopteris fragilis</i>	0.015	5.56	0.01
<i>Dactylis glomerata</i>	0.015	5.56	0.01
<i>Draba</i> spp.	0.015	5.56	0.01
<i>Elymus</i> spp.	0.015	5.56	0.01
<i>Elymus virginicus</i>	0.015	5.56	0.01
<i>Erigeron speciosus</i>	0.015	5.56	0.01
<i>Heuchera</i> spp.	0.015	5.56	0.01
<i>Lepidium densiflorum</i>	0.015	5.56	0.01
<i>Maianthemum stellatum</i>	0.015	5.56	0.01
<i>Melilotus albus</i>	0.015	5.56	0.01
<i>Melilotus officinale</i>	0.015	5.56	0.01
<i>Noccaea montana</i>	0.015	5.56	0.01
<i>Oligosporus dracunculus</i> ssp. <i>glaucus</i>	0.015	5.56	0.01
<i>Oreocarya virgata</i>	0.015	5.56	0.01
<i>Oxybaphus hirsutus</i>	0.015	5.56	0.01
<i>Padus virginiana</i> ssp. <i>melanocarpa</i>	0.015	5.56	0.01
<i>Paronychia jamesii</i>	0.015	5.56	0.01
<i>Penstemon</i> spp.	0.015	5.56	0.01
<i>Poa fendleriana</i>	0.015	5.56	0.01
<i>Podospermum laciniatum</i>	0.015	5.56	0.01
<i>Pterogonum alatum</i>	0.015	5.56	0.01
<i>Rumex triangulivalvis</i>	0.015	5.56	0.01
<i>Schedonnardus paniculatus</i>	0.015	5.56	0.01
<i>Schizachyrium scoparium</i>	0.015	5.56	0.01
<i>Senecio integerrimus</i>	0.015	5.56	0.01
<i>Solanum</i> spp.	0.015	5.56	0.01
<i>Solanum triflorum</i>	0.015	5.56	0.01
<i>Solidago missouriensis</i>	0.015	5.56	0.01
<i>Teloxys botrys</i>	0.015	5.56	0.01
<i>Tithymalus peplus</i>	0.015	5.56	0.01
<i>Triticum aestivum</i>	0.015	5.56	0.01
Unknown forb	0.015	5.56	0.01
<i>Urtica gracilis</i> ssp. <i>gracilis</i>	0.015	5.56	0.01
<i>Vaccinium myrtillus</i> ssp. <i>oreophilum</i>	0.015	5.56	0.01
<i>Yucca glauca</i>	0.015	5.56	0.01

\*Relative Importance = Importance/ Maximum Importance Value in Data Set, Importance = Constancy X Average Cover-All

\*\*Average Cover-All = Sum of first and additional hits.

\*\*\*Indicator Species in TWINSpan Classification

The species with the greatest increase or decrease in importance in 2007.

**Increasers**

	2007 Relative Importance	2002 Relative Importance	Change
Litter	100.00	61.88	38.12
<b>TOTAL INTRO. ANN. GRASSES</b>	27.66	2.40	25.26
<i>Anisantha tectorum</i>	25.88	2.37	23.51
<b>TOTAL NATIVE SHRUBS</b>	30.73	8.45	22.28
<b>TOTAL NATIVE PERENNIAL FORBS</b>	47.18	26.37	20.81
<i>Ceanothus fendleri</i>	22.25	5.10	17.15
Standing dead	14.87	1.74	13.13
<i>Heterotheca foliosa</i>	6.44	0.20	6.24
<i>Artemisia frigida</i>	6.56	0.86	5.70
<i>Achillea lanulosa</i>	3.36	0.09	3.26
<i>Artemisia ludoviciana</i>	4.44	1.24	3.20
<i>Solidago simplex</i> var. <i>simplex</i>	2.64	0.36	2.28
<i>Poa agassizensis</i>	1.53	0.04	1.49
<b>TOTAL NATIVE TREES</b>	4.34	2.92	1.42
<i>Penstemon virens</i>	2.74	1.52	1.22
<i>Cercocarpus montanus</i>	1.26	0.16	1.10
<i>Heterotheca villosa</i>	1.46	0.42	1.04
<i>Elymus elymoides</i>	1.26	0.25	1.01
<i>Pseudotsuga menziesii</i>	1.15	0.47	0.68
<i>Aster porteri</i>	0.82	0.17	0.65
<i>Machaeranthera bigelovii</i>	0.63	0.01	0.62
<i>Ribes cereum</i>	0.68	0.19	0.49
<i>Eriogonum umbellatum</i> var. <i>umbellatum</i>	0.47	0.01	0.46
<i>Rubus idaeus</i> ssp. <i>melanolasius</i>	0.61	0.24	0.36
<i>Koeleria macrantha</i>	0.32	0.01	0.31
<i>Muhlenbergia montana</i>	0.34	0.05	0.29
<i>Acosta diffusa</i>	0.19	0.00	0.19
<i>Alyssum minus</i>	0.48	0.31	0.17
<i>Tragopogon dubius</i> ssp. <i>major</i>	0.13	0.00	0.13
<i>Cerastium strictum</i>	0.12	0.00	0.11
<i>Hesperostipa comata</i>	0.15	0.03	0.11
<b>TOTAL NATIVE PERENNIAL GRASSES (c)</b>	23.90	23.81	0.09
<i>Elymus lanceolatus</i> fm. <i>albicans</i>	0.40	0.33	0.07
<i>Populus tremuloides</i>	0.54	0.47	0.07
<i>Bahia dissecta</i>	0.06	0.00	0.06
<i>Bromopsis lanatipes</i>	0.05	0.00	0.05

**Decreasers**

	Relative Importance	Relative Importance	Change
<i>Bare soil</i>	48.11	100.00	-51.89
<i>Pinus ponderosa</i> ssp. <i>scopulorum</i> [dead]	0.21	9.58	-9.37
<b>TOTAL NATIVE ANN. &amp; BIEN. FORBS</b>	1.55	8.70	-7.15
<i>Phacelia heterophylla</i>	0.48	5.10	-4.61
<b>TOTAL INTRO. ANN. &amp; BIEN. FORBS</b>	7.14	9.91	-2.77
<i>Chenopodium simplex</i>	0.00	2.65	-2.65
<i>Sisymbrium altissimum</i>	0.09	2.69	-2.60
<i>Carex pensylvanica</i> ssp. <i>heliophila</i>	4.15	6.31	-2.16
<i>Ceratochloa carinata</i>	0.00	1.76	-1.76
<i>Elymus trachycaulus</i>	0.52	1.85	-1.33
<i>Pseudotsuga menziesii</i> [dead]	3.16	4.11	-0.96
<b>TOTAL INTRO. PERENNIAL GRASSES (c)</b>	0.14	0.94	-0.80
<i>Chenopodium leptophyllum</i>	0.00	0.78	-0.78
<b>TOTAL INTRO. PERENNIAL FORBS</b>	0.38	1.09	-0.72
<i>Grindelia squarrosa</i>	0.12	0.69	-0.57
<i>Epilobium brachycarpum</i>	0.00	0.56	-0.56
<i>Apocynum androsaemifolium</i>	0.04	0.60	-0.56
<i>Verbascum thapsus</i>	2.95	3.48	-0.54
<i>Astragalus miser</i> var. <i>oblongifolius</i>	0.00	0.52	-0.52
<i>Breea arvensis</i>	0.17	0.61	-0.44
<i>Geranium caespitosum</i> ssp. <i>caespitosum</i>	0.81	1.21	-0.41
<i>Carex</i> spp.	0.67	1.06	-0.39
<i>Chondrosium gracile</i>	0.21	0.57	-0.36
<i>Campanula rotundifolia</i>	0.46	0.78	-0.32
<i>Sabina scopulorum</i> [dead]	0.03	0.32	-0.29
<i>Lactuca serriola</i>	0.10	0.38	-0.27
<i>Silene antirrhina</i>	0.00	0.24	-0.24
<i>Erysimum capitatum</i>	0.05	0.29	-0.24
<i>Helianthus pumilus</i>	0.43	0.67	-0.23
<b>TOTAL NATIVE PERENNIAL GRASSES (w)</b>	0.78	1.00	-0.22
<i>Gaillardia aristata</i>	0.00	0.22	-0.22
<i>Physocarpus monogynus</i>	0.23	0.44	-0.22
<i>Carduus nutans</i> ssp. <i>macrolepis</i>	0.00	0.11	-0.11
<i>Astragalus laxmannii</i>	0.00	0.11	-0.11
<i>Drymocallis fissa</i>	0.09	0.19	-0.09
<i>Acrolasia dispersa</i>	0.00	0.09	-0.09
<i>Pulsatilla ludoviciana</i>	0.00	0.08	-0.08
<i>Scutellaria brittonii</i>	0.00	0.08	-0.08
<i>Androsace septentrionalis</i>	0.00	0.06	-0.06

**Newly observed, or no longer observed in 2007 transects.**

**NEW SPECIES in 2007**

*Elymus lanceolatus*  
*Festuca arizonica*  
*Microseris nutans*  
*Pinus ponderosa* ssp. *scopulorum*  
*Erigeron divergens*  
*Oenothera villosa* ssp. *strigosa*  
*Cynoglossum officinale*  
*Descurainia sophia*  
*Turritis glabra*  
*Erigeron compositus*  
*Erigeron pinnatisectus*  
*Gastrolychnis drummondii*  
*Heliomeris multiflora*  
*Lesquerella montana*  
*Penstemon strictus*  
*Physalis heterophylla*  
*Scrophularia lanceolata*  
*Senecio spartioides*  
*Viola* spp.  
*Linaria genistifolia* ssp. *dalmatica*  
*Achnatherum lettermanii*  
*Amelanchier alnifolia*  
*Ribes aureum*

**SPECIES OBSERVED IN TRANSECTS IN 2002 BUT NOT OBSERVED IN 2007**

*Corydalis aurea*  
*Triticum aestivum* x *elytrigia elongata*  
*Oenothera caespitosa* ssp. *caespitosa*  
*Dracocephalum parviflorum*  
*Helianthus annuus*  
*Bassia sieversiana*  
*Populus tremuloides* [dead]  
*Liatris punctata*  
*Astragalus agrestis*  
*Oxytropis lambertii*  
*Verbena bracteata*  
Moss  
*Cirsium vulgare*  
*Festuca brachyphylla* ssp. *coloradensis*  
*Oreobatus deliciosus*  
*Astragalus shortianus*  
*Nicotiana attenuata*  
*Acer glabrum*  
*Arnica fulgens*  
*Astragalus* spp.  
*Collomia linearis*  
*Euphorbia* spp.  
*Gaura mollis*  
*Physaria vitulifera*  
*Sporobolus cryptandrus*  
*Achnatherum nelsonii*  
*Asclepias stenophylla*  
*Asclepias viridiflora*  
*Boechera fendleri*  
*Carex* sp. 1  
*Chenopodium foliosum*  
*Chrysothamnus parryi*  
*Cystopteris fragilis*  
*Draba* spp.  
*Elymus* spp.  
*Heuchera* spp.  
*Lepidium densiflorum*  
*Melilotus albus*  
*Melilotus officinale*  
*Noccaea montana*  
*Oreocarya virgata*  
*Oxybaphus hirsutus*  
*Poa fendleriana*  
*Podospermum laciniatum*  
*Schedonnardus paniculatus*  
*Schizachyrium scoparium*  
*Senecio integerrimus*  
*Solanum* spp.  
*Solanum triflorum*  
*Teloxys botrys*  
*Tithymalus peplus*  
*Urtica gracilis* ssp. *gracilis*  
*Vaccinium myrtillus* ssp. *oreophilum*

Appendix 3. Photographs

**Photographs  
Walker Ranch  
Boulder County  
July, 2007**



**Sample start point**



**Sample end point**

**Sample Site 1**



**Sample start point**



**Sample end point**

**Sample Site 2**



**Sample start point**



**Sample end point**

**Sample Site 3**



**Sample start point**



**Sample end point**

**Sample Site 4**



**Sample start point**



**Sample end point**

**Sample Site 5**



**Sample start point**




**Sample end point**

**Sample Site 6**



**Sample start point**



**Sample end point**

**Sample Site 7**



**Sample start point**



**Sample end point**

**Sample Site 8**



**Sample start point**



**Sample end point**

**Sample Site 9**



**Sample start point**



**Sample end point**

**Sample Site 10**



**Sample start point**



**Sample end point**

**Sample Site 11**



**Sample start point**



**Sample end point**

**Sample Site 12**



**Sample start point**



**Sample end point**

**Sample Site 13**



**Sample start point**



**Sample end point**

**Sample Site 14**



**Sample start point**



**Sample end point**

**Sample Site 15**



**Sample start point**



**Sample end point**

**Sample Site 16**



**Sample start point**



**Sample end point**

**Sample Site 17**



**Sample start point**



**Sample end point**

**Sample Site 18**

Vertical lines define Sample groups and horizontal lines define species groups.

**Sample Classification Groups**

111 1 11111  
 875371245968106423  
**E D C B A**

\* This symbol identifies the indicator species defined by TWINSPAN

143	Sisy alti	--1-2249-1-31-421	00000
74	Gail aris	-1111211----1-3--	000010
102	Nico atte	-----11-----1	000010
78	Grin squa	-2-2452-----211-	000011
83	Hete foli	-1--1--5-1-----1	000011
104	Oeno caes	--11121----1-2--	000011
159	Verb thap	-27713111121311131	0001
*	10 Amer lanc	11-----	001000
27	Astr shor	-1-11-----	001000
33	Brom lana	-1-1-----	001000
34	Brom japo	-1--1-----	001000
44	Cerc mont	-5--2-11-----	001000
82	Hesp coma	-21-1-1-----	001000
*	90 Lesq mont	-111-----	001000
101	Muhl mont	-4-----1-----	001000
9	Alys minu	--1-316--1-----	001001
35	Came micr	-----11-----	001001
*	38 Care pens	539566682122--11--	001001
50	Cirs ochr	--2-21-----	001001
63	Elym lanc	-2124-1-2--1----	001001
84	Hete vill	1-112242----11----	001001
88	Lapp redo	----1-12-----	001001
100	Moss	3-----	001001
109	Oxyt lamb	-----12-----	001001
128	Pseu menz	9-----	001001
8	Alli cern	-1-1-----1----	001010
22	Aste port	-1112-121-1-111----	001010
23	Astr agre	---2-----1----	001010
30	Bass siev	--2--111-1-----	001010
62	Elym elym	-122111---3-1----	001010
69	Erio umbe	-1--1111--1-1----	001010
79	Harb trac	--111--2--1-1----	001010
91	Leuc king	242121-11-111-1----	001010
92	Liat punc	-1---2-----1----	001010
106	Oreo deli	-11---1-1-----	001010
110	Pack fend	-1-1-----1----	001010
153	Trag dubi	-11-----1----	001010
160	Verb brac	---2---1----	001010
18	Arte frig	1-115-61---1-1-1-	001011
140	Scut brit	-11-21111---1-1--	001011
70	Erys capi	-2112111-111-12---	001100
86	Koel macr	--11--1-1-1-1----	001100
98	Mert lanc	-111-1-1--1111----	001100
119	Pinu ponD	4-65699-67782-4---	001100
131	Puls ludo	-21---1111-111----	001100
4	Acos diff	---1--1-----	001101
121	Poa comp	---2-----2----	001101
127	Pote hipp	-1-----11-----	001101
155	Trit xely	--11111214133----	001101
2	Achi lanu	1-111--21--1111-1-	001110
132	Ribe cere	1211-11--1-21111--	001110
147	Soli simp	1-2---241111--1--	001110
19	Arte ludo	-41121213112--51--	001111
28	Astr spp	-----1-----1---	010000
43	Cera cari	----3---61877----	010010
65	Elym trac	----4---46795----	010010
*	48 Chon grac	-2--2-1-11442----	010011
115	Pens vire	141112-1132312311-	01010
37	Card nuta	-2--1---1-11-21--	010110
120	Poa agas	--1-----211-1-	010111
116	Phac hete	-16331112663--95-4	011000
25	Astr laxm	-1--2----112--1-1	011001
26	Astr mise	1-11112--1-1122-13	011001
61	Drym fiss	1-1-1111-11-121--2	011001
7	Alet acau	11-----12-1--2--1	011010
67	Epil brac	-1---121111-15111	011011
124	Poly doug	-1---11---1-1--	01110
*	12 Anis tect	-11111266-212-16--	011110
77	Gera caes	-1114151111-1115--	011111
81	Heli pumi	-151211--111--221	100000
1	Acer glab	-1-----1	100001
41	Cean fend	-45533221111131268	100001
133	Rosa arka	-1--11-12-----3	100001
118	Phys mono	-2111--1-11-----5-	100010
137	Sabi scop	-5---21-----5--	100010
32	Bree arve	--6-1---21-1-15-	100011
142	Sile anti	1-21--1-1--111-31-	10010
46	Chen lept	-11112211-1-21421	10011
17	Arni fulg	-----11	101001
60	Drac parv	-----1-----51	101001
99	Mona fist	----1-----2--	101001
*	125 Popu trem	-----2-----645	101001
149	Symp rotu	1-----21-	101001
14	Apoc andr	-----1-1-----78	101010
75	Gali sept	-----1-1-1	101011
113	Pens glab	-----1--11	101011
5	Acro disp	-11--1-----113--	101100
40	Care spp	6-1--21----31-324	101101
47	Chen simp	-111-2-1121113-865	10111
135	Rubu idae	--1--11-111--1141	110000
87	Lact serr	-11---11111-1241	110001
150	Tara offi	-11-----12--111	110010
129	Pseu menD	325-----3-756677	110011
36	Camp rotu	1-212-1-11-1123221	1101
54	Cony cana	-----11---1-	111000
80	Heli annu	-----1-21--2--	111001
126	Popu treD	-----2-----2-	111001
55	Cory aure	-1---2-43112-125	11101
*	11 Andr sept	-----1211-1-11	11110
16	Arct ouva	-----21-1--	11111

0000000000000111  
 000000000111111001  
 01111111100000101

**Sample Classification Groups**

0001111100011  
 001000110110  
 01 0010  
**E D C B A**

Appendix 5. 2002 Species & Site Attribute Data


**Attribute Data for  
Species and Sites  
Walker Ranch  
Boulder County  
July, 2002**

## List of Figures for Appendix 5


Figure 1. <i>Pinus ponderosa</i> ssp. <i>scopulorum</i> (dead) Cover Distribution (2002).	98
Figure 2. <i>Carex pensylvanica</i> ssp. <i>heliophila</i> Cover Distribution (2002) [Indicator for Groups C, D, & E].	99
Figure 3. <i>Ceanothus fendleri</i> Cover Distribution (2002) – a non-preferential species.	100
Figure 4. <i>Phacelia heterophylla</i> Cover Distribution (2002) – a non-preferential species.	101
Figure 5. <i>Pseudotsuga menziesii</i> (dead) Cover Distribution (2002).	102
Figure 6. <i>Verbascum thapsus</i> Cover Distribution (2002) – non-preferential species.	103
Figure 7. <i>Sisymbrium altissimum</i> Cover Distribution (2002).	104
Figure 8. <i>Chenopodium simplex</i> Cover Distribution (2002) – non-preferential species.	105
Figure 9. <i>Anisantha tectorum</i> Cover Distribution (2002) – an indicator for Groups C & D.	106
Figure 10. <i>Elymus trachycaulus</i> Cover Distribution (2002).	107
Figure 11. <i>Ceratochloa carinata</i> Cover Distribution (2002) indicator species for Group B.	108
Figure 12. <i>Penstemon virens</i> Cover Distribution (2002) – non-preferential species.	109
Figure 13. <i>Artemisia ludoviciana</i> Cover Distribution (2002).	110
Figure 14. <i>Geranium caespitosum</i> ssp. <i>caespitosum</i> Cover Distribution (2002).	111
Figure 15. <i>Carex</i> spp. Cover Distribution (2002).	112
Figure 16. <i>Corydalis aurea</i> Cover Distribution (2002).	113
Figure 17. <i>Artemisia frigida</i> Cover Distribution (2002).	114
Figure 18. <i>Campanula rotundifolia</i> Cover Distribution (2002) – non-preferential species.	115
Figure 19. <i>Chenopodium leptophyllum</i> Cover Distribution (2002) – non-preferential species.	116
Figure 20. <i>Grindelia squarrosa</i> Cover Distribution (2002).	117
Figure 21. <i>Triticum aestivum</i> x <i>Elytrigia elongata</i> Cover Distribution (2002).	118
Figure 22. <i>Helianthus pumilus</i> Cover Distribution (2002).	119
Figure 23. <i>Breaa arvensis</i> Cover Distribution (2002).	120
Figure 24. <i>Apocynum androsaemifolium</i> Cover Distribution (2002).	121
Figure 25. <i>Chondrosium gracile</i> Cover Distribution (2002).	122
Figure 26. <i>Epilobium brachycarpum</i> Cover Distribution (2002).	123
Figure 27. <i>Leucopoa kingii</i> Cover Distribution (2002).	124
Figure 28. <i>Astragalus miser</i> var. <i>oblongifolius</i> Cover Distribution (2002) – non-preferential species.	125
Figure 29. <i>Populus tremuloides</i> Cover Distribution (2002) – indicator species for Group A.	126
Figure 30. <i>Physocarpus monogynus</i> Cover Distribution (2002).	127
Figure 31. Distribution of Slope % of samples scaled from 100 to 0.	128
Figure 32. Distribution of “Northerly” Aspect scaled from 100 to 0.	129
Figure 33. Distribution of “Easterly” Aspect scaled from 100 to 0.	130
Figure 34. Distribution of Standing Dead Percent Cover (2002).	131
Figure 35. Distribution of Total Vegetation Percent Cover (2002).	132

# Attribute Figures for 2002


The attribute figures present the cover values for selected species at each sample site. This has been superimposed on the environmental vectors to represent the environmental centers of species distributions. The position of the circles indicates the sample site location in the ordination, and the size of the symbol is proportional to the actual cover value for the selected species. The species that were selected were the 31 most important species based on cover and frequency. Species that were identified as indicator species are noted as such in the Figure title. The species are presented in the same order as found in the Table 2. Douglas fir (*Pseudotsuga menziesii*) was included in the Table 2 list, but occurred only in Sample 18 with a cover value of 59%. Sample 18 was the unburned site outside of the burn perimeter and was so different from all of the other samples that it had to be excluded from the ordination, and is not presented below.




**Figure 1. *Pinus ponderosa* ssp. *scopulorum* (dead) Cover Distribution (2002).**



**Figure 2. *Carex pensylvanica* ssp. *heliophila* Cover Distribution (2002) [Indicator for Groups C, D, & E].**



**Figure 3. Ceanothus fendleri Cover Distribution (2002) – a non-preferential species.**



**Figure 4. *Phacelia heterophylla* Cover Distribution (2002) – a non-preferential species.**




Figure 5. *Pseudotsuga menziesii* (dead) Cover Distribution (2002).




Figure 6. *Verbascum thapsus* Cover Distribution (2002) – non-preferential species.




Figure 7. *Sisymbrium altissimum* Cover Distribution (2002).





Figure 8. Chenopodium simplex Cover Distribution (2002) – non-preferential species.






Figure 10. *Elymus trachycaulus* Cover Distribution (2002).




Figure 11. *Ceratochloa carinata* Cover Distribution (2002) indicator species for Group B.




Figure 12. *Penstemon virens* Cover Distribution (2002) – non-preferential species.




Figure 13. *Artemisia ludoviciana* Cover Distribution (2002).





Figure 14. *Geranium caespitosum* ssp. *caespitosum* Cover Distribution (2002).






Figure 16. *Corydalis aurea* Cover Distribution (2002).




Figure 17. *Artemisia frigida* Cover Distribution (2002).




Figure 18. *Campanula rotundifolia* Cover Distribution (2002) – non-preferential species.




Figure 19. *Chenopodium leptophyllum* Cover Distribution (2002) – non-preferential species.




Figure 20. *Grindelia squarrosa* Cover Distribution (2002).




Figure 21. *Triticum aestivum* x *Elytrigia elongata* Cover Distribution (2002).




Figure 22. *Helianthus pumilus* Cover Distribution (2002).




Figure 23. *Brea arvensis* Cover Distribution (2002).




Figure 24. *Apocynum androsaemifolium* Cover Distribution (2002).




Figure 25. *Chondrosium gracile* Cover Distribution (2002).




Figure 26. *Epilobium brachycarpum* Cover Distribution (2002).




Figure 27. *Leucopoa kingii* Cover Distribution (2002).




Figure 28. *Astragalus miser* var. *oblongifolius* Cover Distribution (2002) – non-preferential species.




Figure 29. *Populus tremuloides* Cover Distribution (2002) – indicator species for Group A.




Figure 30. *Physocarpus monogynus* Cover Distribution (2002).




Figure 31. Distribution of Slope % of samples scaled from 100 to 0.




Figure 32. Distribution of “Northerly” Aspect scaled from 100 to 0.




Figure 33. Distribution of "Easterly" Aspect scaled from 100 to 0.




Figure 34. Distribution of Standing Dead Percent Cover (2002).





Figure 35. Distribution of Total Vegetation Percent Cover (2002).

## Appendix 6. Computer files and GPS Sample Coordinates

File inventory:

All of the project related files have been included in a CD. The directory structure is as follows:



The following is an explanation of the contents of the subdirectories:

**AutoCAD** – Contains files that were used to construct the report graphics that included sample locations and treatment and burn areas. The CADImages subdirectory includes the aerial photography and USGS images that were used in the AutoCAD files.

**Classification&Ordination** – Contains input and output files from TWINSPAN and CANOCO. The subdirectory Descrim contains the input and output files from DESCRIM.

**FieldData** – Contains the original vegetation and site data and summary information in EXCEL tables. There are two subdirectories for the years 2002 and 2007. The site data are included as worksheets in the Excel spreadsheets.

**FinalReport** - The final reports for both 2002 and 2007 are included as both a Microsoft Word 2003 documents (WalkerRept200X.doc) and as an Adobe .pdf file (WalkerRept200X.pdf).

**Photographs** – Contains all of the photographs for the sample sites. There are two subdirectories for the years 2002 and 2007.

Ecotone - 2002/2007												
BCOS Walker Ranch Sample Coordinates in UTM NAD 27 meters.												
b = endpoint location												
Sample	Easting	Northing	Elevation meters	Elevation feet	Sampler	Orientation	Aspect	Slope°	Slope%	DATE	DATE	COMMENT 2007
01W	469,880.84	4,422,661.00	2,291.7	7,516.9	PM	290	4	10	17.60	7/23/2007	7/18/2002	
01Wb	469,841.83	4,422,693.79	2,284.0	7,491.6								
02W	470,317.47	4,422,385.03	2,221.0	7,284.8	PM	35	154	16	28.70	7/21/2007	7/17/2002	
02Wb	470,354.33	4,422,416.80	2,223.6	7,293.4								
03W	470,581.10	4,422,307.06	2,191.1	7,186.8	PM	175	144	16	28.7	7/22/2007	7/17/2002	
03Wb	470,575.50	4,422,259.17	2,181.3	7,154.7								
04W	470,487.83	4,421,647.76	2,187.5	7,175.2	PM	210	114	15	26.8	7/21/2007	7/18/2002	
04Wb	470,456.64	4,421,609.97	2,189.9	7,183.0								
05W	470,584.96	4,421,466.51	2,192.2	7,190.3	PM	228	48	16	28.7	7/22/2007	7/18/2002	
05Wb	470,545.09	4,421,439.67	2,208.8	7,244.8								
06W	470,616.60	4,422,709.63	2,193.8	7,195.8	PM	360	274	22	40.4	7/21/2007	7/19/2002	White fiberglass at start point
06Wb	470,627.83	4,422,758.68	2,198.8	7,212.1								Carsonite post at end point
07W	470,416.57	4,422,424.97	2,222.4	7,289.6	PM	278	178	10	17.6	7/22/2007	7/17/2002	
07Wb	470,368.43	4,422,440.47	2,221.6	7,286.8								
08W	470,983.61	4,422,261.87	2,254.5	7,394.7	PM	360	330	18	32.5	7/24/2007	7/16/2002	
08Wb	470,982.53	4,422,307.95	2,234.3	7,328.4								
09W	470986.76 (47096)	4422349.39 (4422356)	2,215.1	7,265.7	PM	240	320	18	32.5	7/24/2007	7/16/2002	Start was moved about 3m east due to new trail. Transect crosses trail at meter 2.5-7.5. Took new GPS at start and end.
09Wb	470,938.64	4,422,341.08	2,210.8	7,251.3								
10W	470,376.61	4,422,757.10	2,227.1	7,304.7	PM	30	80	6	10.5	7/23/2007	7/18/2002	The old GPS coordinates were off, tried to get new data
10Wb	470,408.97	4,422,792.93	2,222.5	7,289.7								
11W	471,153.67	4,422,071.15	2,204.4	7,230.3	PM	25	124	18	32.5	7/24/2007	7/19/2002	
11Wb	471,177.92	4,422,113.92	2,210.8	7,251.3								
12W	470,050.63	4,422,210.84	2,306.2	7,564.3	PM	147	72	20	36.4	7/21/2007	7/17/2002	Could not find end point, placed new white fiberglass pole
12Wb	470,067.78	4,422,165.66	2,320.7	7,611.8								
13W	470,261.88	4,422,259.07	2,222.5	7,289.7	PM	158	52	24	44.5	7/21/2007	7/17/2002	Could not find end point, placed new white fiberglass pole.
13Wb	470,274.09	4,422,214.13	2,241.5	7,352.1								
14W	470,293.31	4,422,682.78	2,227.9	7,307.4	PM	248	70	14	24.9	7/21/2007	7/18/2002	
14Wb	470,242.65	4,422,681.22	2,236.1	7,334.5								
15W	472,036.59	4,421,527.00	2,120.5	6,955.2	PM	124	54	22	40.4	7/23/2007	7/19/2002	Could not find start or end markers. Set new white fiberglass poles and GPS'd the new locations.
15Wb	472,071.82	4,421,492.05	2,108.8	6,916.7								
16W	471,447.06	4,421,997.79	2,256.4	7,400.8	PM	252	298	14	24.9	7/23/2007	7/16/2002	
16Wb	471,400.66	4,421,996.49	2,257.0	7,402.9								
17W	471,402.22	4,421,877.77	2,250.8	7,382.5	PM	200	274	20	36.4	7/24/2007	7/19/2002	Transect crosses trail at meters 36-43.
17Wb	471,375.28	4,421,833.91	2,242.9	7,356.8								
18W	471,595.57	4,422,000.41	2,256.7	7,402.1	PM	315	30	18	32.5	7/21/2007	7/19/2002	GPS data were off. Recollected GPS
18Wb	471,578.28	4,422,047.33	2,247.7	7,372.5								